



UNITED ARAB EMIRATES UNIVERSITY



# **2023 UAEU-AUA International Workshop on Delay Differential Equations Applications to Immunology and Infectious Diseases (DDEs-AIIDs)**

from 23–24 Nov., 2023, UNITED ARAB EMIRATES UNIVERSITY

College of Science  
Department of Mathematical Sciences  
UAE  
23–24 November, 2023

## **Book of Abstracts**



## **Contents**

<b>1 Organizing Committee</b>	<b>1</b>
<b>2 Workshop Overview</b>	<b>2</b>
<b>3 Keynote Speakers</b>	<b>2</b>
<b>4 Invited Speakers</b>	<b>2</b>
<b>5 Abstracts</b>	<b>4</b>
<b>6 List of Participants</b>	<b>89</b>

## **1 Organizing Committee**

1. Prof. Fathalla A. Rihan (Chair, UAEU), Email: frihan@uaeu.ac.ae
2. Dr. Abdessamad Tridane
3. Dr. Ardak Kashkynbayev
4. Dr. Khalid Muhammad

### **Logistic Committee:**

- |                               |                         |
|-------------------------------|-------------------------|
| 1. Prof. Mohamed Syam         | 5. Dr Bilal Rather      |
| 2. Dr. Hebatallah J. Alsakaji | 6. Dr. Mohammed Abuomar |
| 3. Dr. Jianhua Gong           | 7. Ms. Ghilmana Sarmad  |
| 4. Dr. Mohammed Imran         | 8. Ms. Fareeha Jamal    |

### **Scientific Committee:**

- |  |                                   |
|--|-----------------------------------|
| 1. Prof. Fathalla A. Rihan (UAE)       | 7. Prof. Ephraim Agyingi (USA)    |
| 2. Dr. Ardak Kashkynbayev (Kazakhstan) | 8. Prof. Radouane Yafia (Morocco) |
| 3. Prof. G. Bocharov (Russia)          | 9. Prof. Nasser Sweilam (Egypt)   |
| 4. Prof. Andreas Meyerhans (Spain)     | 10. Prof. Cemil Tunc (Turkey)     |
| 5. Prof. Yang Kuang (USA)              | 11. Prof. M-Naim Anwar (Egypt)    |
| 6. Prof Hongjiong Tian (China)         | 12. Dr. R. Rakkiyappan (India)    |

## **2 Workshop Overview**

A growing interest has been shown in delay differential equations (DDEs) due to their important applications in biology, medicine, immunology, and epidemiology. Using mathematical models and DDEs of infectious diseases, we can study the pathogenesis of disease processes, predict outbreaks, and minimize outbreak severity and frequency. This focused workshop DDEs-AIID will bring together mathematicians and life science scholars at AUA universities, as well as outside of AUA universities, to meet, share, and discuss research interested in mathematical modeling with delay differential equations and their applications to immunology and infectious diseases. There will be a round table discussion on open problems and questions in the context of infectious diseases, cancer, and chronic diseases, as well as future directions in multidisciplinary research. Potential topics include but are not limited to the following i) DDEs mathematical analysis and biological interpretation, ii) Models of host-pathogen interactions, iii) Models of cancer immune response, iv) Models of immunological regulation, v) Therapy and vaccination models, vi) Models of multiscale immunoepidemiology, vii) Optimal control with therapy, viii) Immune response, signaling pathways, and regulatory networks

The scientific program of the workshop consists of 50-minute plenary lectures, 30-minute invited lectures and 20-minute contributed talks.

The selected papers presented will be published in the Special Issue "Delay Differential Equations and Applications to Immunology and Infectious Diseases" of "Mathematical Biosciences and Engineering", published by AIMS.

## **3 Keynote Speakers**

1. Prof. Gennady Bocharov, Marchuk Institute of Numerical Mathematics, Russian Academy of Sciences, Moscow, Russia;
2. Prof. Andreas Meyerhans, Infection Biology Laboratory Department of Medicine and Life Sciences, Universitat Pompeu Fabra, Barcelona, Spain;
3. Prof. Vitaly Volpert, Institut Camille Jordan, UMR 5208 CNRS, University Lyon 1, 69622 Villeurbanne, France.

## **4 Invited Speakers**

1. Prof. Radouane, Yafia, Morocco: Ibn Tofail University
2. Prof. Abdulla Azamov, Uzbekistan: Institute of Mathematics
3. Dr. Ardak KASHKYNBAYEV, Kazakhstan: Nazarbayev University
4. Prof. Jiaxu Li, USA: Department of Mathematics, University of Louisville

### **Acknowledgement**

We are grateful for the support of Department of Mathematical Sciences (UAEU), Asian Universities Alliance (AUA) for support and helping make this workshop possible. . . . Fathalla A. Rihan

## 5 Abstracts

# Understanding the pathogenesis of virus infections using mathematical models based on delay differential equations

G. Bocharov<sup>1,2,3</sup>

<sup>1</sup> Marchuk Institute of Numerical Mathematics, Russian Academy of Sciences (INM RAS), Moscow

<sup>2</sup> Moscow Center for Fundamental and Applied Mathematics at INM RAS, Moscow

<sup>3</sup> Sechenov First Moscow State Medical University, Moscow

Email: bocharov@m.inm.ras.ru

### Abstract

Mathematical immunology is a rapidly developing highly demanded field of applied mathematics. It aims to bring the rigor of mathematics and the power of computer technologies to describe, analyze and predict the dynamic complexities of the immune system. The immune system functions to maintain the antigenic homeostasis of the host organism. Virus infections induce a set of antiviral immune responses with the outcome depending on the dynamics of the pathogen-host interaction. We review the structure of representative mathematical models describing processes of multi-physics nature formulated with delay differential equations. We apply the models to studying the pathogenesis of infections such as SARS-CoV-2, HIV-1, HBV and to quantifying the growth kinetic parameters of CFSE labelled lymphocytes. To this end, we apply a range of analytical methods including the sensitivity-, bifurcation-, and dynamic uncertainty analyses. The models provide core building blocks for developing a novel class of mathematical models and methods for implementation of a multi-physics description and analysis of the functioning of immune system under normal conditions and in the course of infectious diseases.

The study is supported by the Russian Science Foundation (grant number 23-11-00116).

# Pathogenic virus infections: on fate regulation and functional cure strategies

Andreas Meyerhans<sup>1,2</sup>

<sup>1</sup>Infection Biology Laboratory, Department of Medicine and Life Sciences (MELIS), Universitat Pompeu Fabra, Barcelona, Spain.

<sup>2</sup> Instituci Catalana de Recerca i Estudis Avanats (ICREA), Barcelona, Spain.

Email : andreas.meyerhans@upf.edu

## Abstract

The fates of pathogenic virus infections are determined by the dynamic interplay between the expanding virus and the concomitantly induced immune response in the infected host. Depending on the temporal virus-host relationship, infections are either acute or chronic. In humans, acute infections are usually resolved within few weeks. In contrast, chronic infections are not resolved and, instead, develop when innate and adaptive immune responses are not sufficient to eliminate the invading virus during the primary infection phase. This presentation will address 2 fundamental questions of virus infections: (1) Which immunological features of the infected host organism determine the acute or chronic outcome of a virus infection? (2) Can chronic persistent virus infections be functionally cured by immunotherapeutic interventions? Experiments using the acute and chronic lymphocytic choriomeningitis virus (LCMV) infection model of mice will be presented and discussed. They provide clues about immune system perturbation and functioning that may be of general importance also for other infections like those with the human immunodeficiency virus (HIV).

This work is supported by grants from the Spanish Ministry of Science and Innovation grant No. PRPPID2022-141395OB-I00, la Caixa Foundation (HR17-00199), the Unidad de Excelencia Mara de Maeztu, funded by the AEI (CEX2018-000792-M), and the Russian Science Foundation (grant number 23-11-00116).

## References

- [1] Bocharov G, Argilaguet J, Meyerhans A. Understanding Experimental LCMV Infection of Mice: The Role of Mathematical Models. *J Immunol Res.* 2015, 739706. doi: 10.1155/2015/739706.
- [2] Argilaguet J, Pedragosa M, Esteve-Codina A, Riera G, Vidal E, Peligero-Cruz C, Casella V, Andreu D, Kaisho T, Bocharov G, Ludwig B, Heath S, Meyerhans A. Systems analysis reveals complex biological processes during virus infection fate decisions. *Genome Res.* 2019 29(6):907-919. doi: 10.1101/gr.241372.118.

## Mathematical modelling of respiratory viral infections

V. Volpert <sup>\*1</sup>

<sup>1</sup>Institut Camille Jordan, UMR 5208 CNRS, University Lyon 1, 69622 Villeurbanne, France

Email : volpert@math.univ-lyon1.fr

### **Abstract**

In this lecture we will present an overview of recent works on mathematical modelling of respiratory viral infections. We will begin with the investigation of infection progression in cell cultures and in tissues of human body. We will determine viral load and infection spreading speed and we will apply these results to evaluate infectivity and severity of symptoms for different variants of the SARS-CoV-2 infection. Next, will take into account mucus motion in the lung airways and will show how viral infection can lead to mucus accumulation and airway obstruction leading to a life-threatening disease progression.

# Qualitative and Quantitative Features of Delay Differential Equations and Applications to Immunology

Fathalla A. Rihan <sup>1</sup>

<sup>1</sup> Department of Mathematical Sciences, College of Science, United Arab Emirates University, Al Ain, 15551, UAE

Email: frihan@uaeu.ac.ae

## Abstract

Recently much attention has been given to mathematical modeling of real-life phenomena using differential equations with memory, such as delay differential equations (DDEs). This is due to the fact that introduction memory terms in a differential model significantly increases the complexity of the model. Such a class of DDEs is widely used for analysis and predictions in various areas of life sciences and modern topics in population dynamics, computer science, epidemiology, immunology, physiology, and neural networks. In this talk, we provide a wide range of delay differential models that have a richer mathematical framework (compared with ODEs) for the analysis of biosystems. Qualitative and quantitative features of DDEs are discussed. Some numerical simulations are also provided with tumour-immune interactions to show the effectiveness of the theoretical results.

## References

- [1] Rihan F.A., Delay differential equations and applications to biology, Springer (SP) (2021).



# Dynamics of A Delayed Spatiotemporal HBV Infection Model with Capsids, CTL Immune Response and General Incidence Function

Abdessamad Tridane<sup>1</sup>

<sup>1</sup> Department of Mathematical Sciences, College of Science, United Arab Emirates University, Al Ain, UAE

Email: a-tridane@uaeu.ac.ae

## Abstract

This study explores the integrated effects of diffusion and CTL immune response activation delay on the in vivo dynamics of a generalized hepatitis B virus (HBV) infection model in terms of both mathematical analysis and numerical simulations. The considered model involves five compartments such as uninfected and infected hepatocytes, HBV DNA-containing capsids, viruses, and CTL cells. We model the infection process by a general incidence function with some biologically feasible assumptions and, as a result, it includes a wide variety of popular incidence functions utilized in the existing literature. We then establish the well-posedness of the considered model in terms of the existence, uniqueness, non-negativity, and boundedness of the solutions. Further, three different steady states such as disease-free, immune-free and infection steady states are identified for the model under threshold conditions depending upon the reproduction numbers  $\mathcal{R}_i$  and  $\mathcal{R}_{CTL}$ . With the help of Lyapunov functions and LaSalle's invariance principle, we prove that both the disease-free and immune-free steady states are globally asymptotically stable under suitable conditions. On the other hand, infection steady state with CTL immune response exhibits the Hopf bifurcation with respect to the delay parameter. Finally, some corroborative numerical simulations are presented with a particular choice of incidence function.

# Applications of Delay Differential Equations in Medicine

Fathalla A. Rihan<sup>1</sup> and Adel Hashish\*<sup>2</sup>

<sup>1</sup> Department of Mathematical Sciences, College of Science, UAE University, Al-Ain, UAE

<sup>2</sup>Department of Physics, College of Science, UAE University, Al-Ain, UAE

Email : ahashish@uaeu.ac.ae

## Abstract

**Motivation and Aims:** Recently much attention has been given to mathematical modeling of real-life phenomena using differential equations with memory, such as delay differential equations (DDEs). This is due to the fact that introduction memory terms in a differential model significantly increases the complexity of the model. Such a class of DDEs is widely used for analysis and predictions in various areas of life sciences and modern topics in population dynamics, computer science, epidemiology, immunology, physiology, and neural networks. In this paper, we provide a wide range of delay differential models with a richer mathematical framework (compared with ODEs) for analyzing biosystems. Qualitative and quantitative features of DDEs are discussed. Some numerical simulations are also provided to show the effectiveness of the theoretical results.

**Results and Conclusions:** 1. DDEs of real-life phenomena have potentially more interesting dynamics than equations without memory; 2. The time delay (memory) is incorporated into systems to describe resource regeneration times, maturation periods, reaction times, feeding times, gestation periods, etc.; 3. Studying the qualitative behaviour of DDEs is essential for ensuring safe applications or real problems; 4. The presence of memory in the model enriches the dynamics of the model: It can stabilize or destabilize the system. 5. Mono-implicit continuous RK schemes are stable and suitable for stiff and nonstiff DDEs. 6. Sensitivity functions clearly demonstrate the measure of the importance of the input parameters. 7. Sensitivity functions enable one to assess the relevant time intervals for the identification of specific parameters.

## References

- [1] Rihan, F.A., *Delay Differential Equations and Applications to Biology*, Springer (SP), 2021;

# Global behavior in differential delay models of population dynamics and physiology

Anatoli Ivanov<sup>1</sup>

<sup>1</sup> Department of Mathematics, Pennsylvania State University, USA

Email: aivanov@psu.edu

## Abstract

Dynamics of solutions in simple form scalar delay differential equations are studied. The first class includes the Mackey-Glass type equations modeling several distinct processes such as population fluctuations, mechanisms of blood cell production, some others [2, 3, 4]. The second equation is recently proposed as a mathematical model of megakaryopoiesis (platelet production) [1]. Sufficient conditions are derived when the unique positive equilibrium is globally asymptotically stable. The stability conditions are given in terms of induced interval maps, one set being delay independent criteria and another one involving the delay. The existence of periodic solutions slowly oscillating about the equilibrium is also established. The periodic oscillations in the models exist when the positive equilibrium is linearly unstable. The proof of existence can be done by several methods, in particular by following the established ejective fixed point techniques with necessary modifications due to the specific form of the equations.

## References

- [1] L. Boullu, M. Adimy, F. Crauste, and L. Pujo-Menjouet, *Oscillation and asymptotic convergence for a delay differential equation modeling platelet production*, *Discrete and Continuous Dynamical Systems, Ser. B* 24:6, 2417–2442, 2019.
- [2] Y. Kuang, *Delay Differential Equations with Applications in Population Dynamics*. Academic Press Inc. 1993, 398 pp. Series: Mathematics in Science and Engineering, Vol. 191.
- [3] M.C. Mackey, *Unified hypothesis of the origin of aplastic anaemia and periodic hematopoiesis*. *Blood*, 51, 941-956, 1978.
- [4] L. Pujo-Menjouet, *Blood cell dynamics: half a century of modelling*, *Math. Model. Nat. Phenom.*, 11:1, 92–115, 2016.

# Crossover Dynamics of Hybrid Fractional Order Monkeypox Disease Model with Time Delay: Numerical Simulations

N. H. Sweilam\*<sup>1</sup> and , S. M. Al-Mekhlafi<sup>2</sup>

<sup>1</sup> Mathematics Department, Faculty of Science, Cairo University, Giza, Egypt

<sup>2</sup> Mathematics Department, Faculty of Education, Sana'a University, Yemen

Email : nsweilam@sci.cu.edu.eg

## Abstract

Monkeypox is a rare viral zoonic illness caused by the monkeypox virus. It was first, identified in 1958. The virus-causing monkeypox comes from the same family of viruses as the variola virus, which causes smallpox. It was first reported in central Africa in 1970. Its symptoms are fever, rash, and lymphadenopathy. In extreme cases, carriers can have pneumonitis, encephalitis, sight-threatening keratitis, and secondary bacterial infections. Children, pregnant women, and immunocompromised persons may be at risk of more severe disease with monkeypox. In this talk, a mathematical model of monkeypox disease is extended using the fractional stochastic-hybrid fractional order deterministic piecewise derivatives. A parameter  $\xi$  is presented in order to be consistent with the physical model problem. The positivity, boundedness, and existence of the solutions for the proposed model are discussed. New numerical algorithms are improved to study the proposed model. These methods are the nonstandard Caputo proportional constant Adams-Bashfourth sixth step method and the nonstandard Euler Maruyama technique to solve the fractional stochastic model. Several numerical tests demonstrate the efficiency of the method and support the theoretical results. The new data simulation and various solution approaches are the novelty of this research. This work aims to explore the transmission pattern of a monkeypox outbreak in the United States.

# Deterministic and stochastic SIRC epidemic model with time-delay for COVID-19

Mohamed-Naim Anwar\*<sup>1</sup>, Hebatallah J. Alsakaji <sup>2</sup> and Fathalla A. Rihan <sup>2</sup>

<sup>1</sup>Pharos University, Alexandria, Egypt

<sup>2</sup> UAE University, AlAin, UAE

mohamed.naeem@pua.edu.eg

## Abstract

An epidemic model based on deterministic and stochastic SIRC has been proposed for COVID-19, taking into account cross-immune classes and time delays in transmission terms. Under some criteria, infection-free and endemic steady states are asymptotically stable. Using the next-generation matrix method, the reproduction number  $\mathcal{R}_0$  for the deterministic model is calculated according to contact rate, recovery rate, and other parameters. Positive global solutions exist and are unique for stochastic analysis. Stochastic models have a smaller reproduction number than deterministic models of the same type  $\mathbb{R}_0$ . We obtain conditions for the extinction of the disease by using the stochastic Lyapunov functional. It appears that white noise and time delay play a significant role in controlling the spread of the virus.

## References

- [1] Rihan F.A., Anwar M., Qualitative analysis of delayed SIR epidemic model with a saturated incidence rate, *International Journal of Differential Equations*, 408637,2012.
- [2] Rihan F.A., Alsakaji H., Rajivganthi C., Stochastic SIRC epidemic model with time-delay for COVID-19, *Advances in difference equations*, 1-20,2020.
- [3] Rihan, F.A., K. Udhayakumar, N. Spinelli, M-N. Anwar, A. Q.M. Khaliq, Stability and Bifurcation Analysis of the Caputo Fractional-order Asymptomatic COVID-19 Model with Multiple Time-delays, *International Journal of Bifurcation and Chaos*, Vol. 33, No. 02, (2023)
- [4] Rihan F.A., *Delay differential equations and applications to biology*, Springer (2021).

# Exploring the Influence of Fear on the Progression of an Infectious Disease in a Delayed Spatiotemporal Epidemic Model Featuring Both Local and Nonlocal Dispersion

Ghilmana Sarmad<sup>1</sup>

<sup>1</sup> Department of Mathematics, College of Science, United Arab Emirates University

Email : 202190243@uaeu.ac.ae

## Abstract

The paper investigates an epidemic model known as the Susceptible-Protected-Infected-Recovered (SPIR). This model incorporates a mixed diffusion framework, comprising both local and nonlocal diffusions, to capture the influence of fear among the population when it comes to infection transmission. We establish the mathematical well-posedness of this model, which entails demonstrating the existence, positivity, and uniqueness of its solution. One key aspect of our analysis involves deriving a variational expression for the basic reproduction number, denoted as  $\mathfrak{R}_0$ . This parameter serves as a crucial threshold for the epidemic dynamics. Specifically, when  $\mathfrak{R}_0$  is less than 1, we establish that the epidemic will eventually die out, signifying the global asymptotic stability of the infection-free equilibrium state. However, when  $\mathfrak{R}_0$  exceeds 1, we demonstrate that the solution will persist uniformly, and an endemic equilibrium state emerges. To ascertain the global stability of this endemic equilibrium, we employ Lyapunov function techniques. We consider two distinct scenarios: one where the diffusion coefficient for susceptible populations is set to zero, and another where the diffusion coefficient for infected populations is set to zero. Furthermore, we conduct a comparative analysis with the classical SIR epidemic model to determine the necessary protective measures required to bring the disease under control. This involves reducing  $\mathfrak{R}_0$  below one, which can be achieved through the implementation of appropriate protection strategies.

# Time Lag Effect of Prey Predator Model with Allee Effect

Kalyan Das\*<sup>1</sup>

<sup>1</sup>Department of Basic and Applied Sciences National Institute of Food Technology Entrepreneurship and Management, HSIIDC Industrial Estate, Kundli, Haryana, India

daskalyan27@gmail.com

## Abstract

We have studied a predator-prey system with time lag impact for intra-specific interaction and a substantial Allee effect on prey population growth. We examined the stability of coexisting equilibrium points using the discrete time delay as a bifurcation parameter and showed the existence Hopf-bifurcation when the discrete delay crosses a lumping parameter value. The normal form technique and centre manifold theory have been used to study the aspect of direction and stability of the periodic Hopf-bifurcating solution. All the analytical results are compared with numerical simulations for biological significance.

**Keyword:** Allee-effect, Stability, Hopf-bifurcation, Intra-specific competition.

# A fractional-order delay differential tumor-immune model: external treatment and optimal control strategies

Fathalla A. Rihan<sup>1</sup>, and K. Udhayakumar\*<sup>2</sup>

<sup>1</sup>Department of Mathematical Sciences, College of science, United Arab Emirates University, Al-Ain, 15551, UAE

<sup>2</sup> Zayed University, Abu Dhabi, UAE

Email : udhai512@gmail.com

## Abstract

In the context of external treatment strategies, this paper aims to present a mathematical model of the tumor-immune system. The governing equations of the model are fractional-order delay differential equations, which incorporate a control variable. The inhibition of transforming growth factor beta by itself to provide restricted clinical benefits. However, when given combined with external treatments, it has the potential to significantly enhance the anti-tumor response of the immune system. The purpose of this study is to employ optimal control theory in order to minimize the cost associated with external treatment strategies and decrease the burden of tumor cells. It has been shown that the model's solutions are non-negative and that the existence of an optimal control. We solve the state system forward in time and the adjoint system backward in time to get a numerical approximation of the optimal control problem. Numerical simulations demonstrate that combination of the transforming growth factor beta and external treatments can significantly reduce the tumor load within a short time frame.



# Robust exponential $H_\infty$ filtering for discrete-time complex-valued neural networks with time-varying delays

SOUNDARARAJAN GANESAN\*<sup>1</sup>, G. NAGAMANI<sup>2</sup> and ARDAK KASHKYNBAYEV<sup>1</sup>

<sup>1</sup>Department of Mathematics, School of Sciences and Humanities, Nazarbayev University, Astana-020000, Kazakhstan.

<sup>2</sup>Department of Mathematics, The Gandhigram Rural Institute (Deemed to be University), Gandhigram-624302, India

Email : soundararajan.ganesan@nu.edu.kz

## Abstract

This paper aims to design a compatible filter for a class of delayed discrete-time complex-valued neural networks (CVNNs) having uncertain weighting parameters subject to the  $H_\infty$  performance measure. For this notion, the complex-valued filter scheme is designed for the proposed uncertain CVNNs with regard to the available output measurements. Following the idea of weighted summation inequalities (WSIs) in [1] and complex-valued reciprocal convex matrix inequality [2], we have improved the new complex-valued WSIs without requiring any extra decision variable to bound the summation terms included in Lyapunov-Krasovskii functional (LKF). By utilizing these newly proposed complex-valued WSI and prescribed LKF, an attempt has been made to design the robust  $H_\infty$  filter such that the filtering error system attains exponential stability with the appropriate filtering gain matrices. Finally, a numerical example is performed to acknowledge the significance and efficiency of the proposed filtering scheme.

## References

- [1] Saravanakumar, R., Stojanovic, S. B., Radosavljevic, D. D., Ahn, C. K., Karimi, H. R. (2019). Finite-time passivity-based stability criteria for delayed discrete-time neural networks via new weighted summation inequalities. *IEEE Transactions on Neural Networks and Learning Systems*, 30(1), 5871.
- [2] Ganesan, S., Gnaneswaran, N. (2020). State-feedback filtering for delayed discrete-time complex-valued neural networks. *IEEE Transactions on Neural Networks and Learning Systems*, 31(11), 47264736

# Optimal Control of Stochastic Michaelis-Menten Kinetics Based Tumour-Immune Interaction with Time Delay

Hebatallah J. Alsakaji <sup>\*1</sup> and Fathalla A. Rihan <sup>1</sup>

<sup>1</sup> Department of Mathematical Sciences, College of Science, United Arab Emirates University, Al Ain, UAE

Email : heba.sakaji@uaeu.ac.ae

## Abstract

Herein, we study an optimal control problem of stochastic model to describe the dynamics of tumour-immune interactions, where the interaction of tumour cells with the immune system is described as Michaelis-Menten. The model includes constant time delay to justify the time needed by the immune system to develop a suitable response after recognizing [1]. We first discuss stability and Hopf bifurcation of the deterministic system [2]. We then explore stochastic stability, and the dynamics of the system in view of environmental fluctuations. Criteria for persistence and sustainability are discussed. Using multiple Lyapunov functions, some sufficient criteria for tumour cell persistence and extinction are obtained. By applying optimal control theory, we seek to optimize the objective function in a random environment and to reduce load of of tumour cells [3]. Optimality conditions and characterization of the control are also discussed. We numerically approximate the solution of the optimal control problem by solving the state system forward and adjoint system backward in time

## References

- [1] Alsakaji H., Stochastic Delay Differential Equations with Applications in Ecology and Epidemics, (2020). Dissertations. 185.
- [2] Rihan F., Alsakaji H., Kundu S. , Mohamed O., Dynamics of a timedelay differential model for tumour-immune interactions with random noise, Alexandria Engineering Journal, 61(12), 11913-11923, 2022.
- [3] Rihan F., Delay differential equations and applications to biology, Springer (2021).

# Turing Bifurcation Induced by Cross-Diffusion and Amplitude Equation in Oncolytic Therapeutic Model

Radouane Yafia<sup>\*1</sup>, Fatiha Najm<sup>1</sup> and M. A. Aziz Alaoui<sup>2</sup>

<sup>1</sup>Department of Mathematics Faculty of Sciences, Ibn Tofail University Campus Universitaire, BP 133, Knitra, Morocco

<sup>2</sup> Normandie Univ, France; ULH, LMAH F-76600 Le Havre; FR-CNRS-3335, ISCN 25 rue Ph. Lebon, 76600 Le Havre, France

radouane.yafia@uit.ac.ma

## Abstract

In this paper, we propose a reaction-diffusion mathematical model augmented with self/cross-diffusion in 2D domain which describes the oncolytic virotherapy treatment of a tumor with its growth following the logistic law. The tumor cells is divided into uninfected and infected cells and the virus transmission is supposed to be in a direct mode (from cell to cell). In the absence of cross-diffusion, we establish well posedness of the problem, nonnegativity and boundedness of solutions, non-existence of positive solutions, local and global stability of the nontrivial steady state and the non-occurrence of Turing instability. In the presence of cross-diffusion, we prove the occurrence of Turing instability by using the cross-diffusion coefficient of infected cells as a parameter. To have an idea about different patterns, we derive the corresponding amplitude equation by using the nonlinear analysis theory. In the end, we perform some numerical simulations to illustrate the obtained theoretical results.

## References

- [1] Komarova, N. L. & Wodarz, D. *ODE models for oncolytic virus dynamics*, J. Theor. Biol. 263, 530543 (2010).
- [2] Smith J., *How to add could*. Acta Mathematica, 268(1), 12012–12312, 2021.
- [3] Najm, F., Yafia, R. & Aziz-Alaoui, M. A. ,*Hopf Bifurcation in Oncolytic Therapeutic Modeling: Viruses as Anti-Tumor Means with Viral Lytic Cycle*, International Journal of Bifurcation and Chaos, 32 (11), 2250171, 2022.
- [4] Wodarz, D. & Komarova, N. , *Computational biology of cancer: Lecture notes and mathematical modeling*, World Scientific, 2008.

# Mathematical Model Integrating Vaccine and Variable Immunity Period in Infectious Disease Dynamics

Motassem Al-arydah\*<sup>1</sup>

<sup>1</sup>Department of Mathematics, Khalifa University, P.O.Box: 127788, Abu Dhabi, UAE

[motassem.alarydah@ku.ac.ae](mailto:motassem.alarydah@ku.ac.ae)

## Abstract

We formulate a mathematical model for an infectious disease incorporating a vaccine and variable immunity duration. Our model integrates the behaviors of susceptible, infected, and recovered individuals, along with the impacts of vaccination and the waning of immunity. We establish a formula for the fundamental reproductive rate and employ it to explore the local asymptotic stability of both the disease-free equilibrium (DFE) and the endemic equilibrium (EE). Our findings reveal that the basic reproduction number is highly sensitive to several factors, most notably the rate of immunity loss. Through time series analysis, we evaluate the models outcomes both with and without vaccination, underscoring the vaccines significance in disease management. Furthermore, we estimate time series outcomes for both constant and variable immunity durations, yielding divergent results. Our conclusions underscore not only the necessity of vaccination but also the critical importance of the vaccines conferred immunity period.

# Improved lung model for predicting aerosol deposition in lung airways

Adel Hashish<sup>\*1,2</sup> and Trevor Williams<sup>2</sup>

<sup>1</sup>Department of Physics, Faculty of Science, UAE University, Al-Ain, P.O. Box: 15551, UAE

<sup>2</sup> Formerly of Applied Electrostatics Research Group, Department of Electrical Engineering, University of Southampton, Highfield, Southampton, SO17 1BJ, UK.

Email : ahashish@uaeu.ac.ae

## Abstract

A computer model, for predicting deposition of therapeutic or pollutant aerosol particles in the human lung, has been improved to calculate deposition in each airway generation during inhalation, exhalation, and the breath-holding pause. Deposition in an airway is still based on aerosol flow between characteristic curves in the solution domain but the calculation can be simplified as the product of local flow rate and time interval has been found to be independent of airway generation number though the separate components are not. The model is versatile as it can handle continuous aerosols or pulsed ones intended to target specific regions where lung disease may exist. Electrostatic deposition mechanisms may be considered as well as conventional ones: sedimentation, diffusion, and impaction, which can be made operative as required. Functional residual capacity and head volume are input parameters as are particle diameter, charge, density, and initial number concentration. Inhalation and exhalation flow rates and durations, as well as the breath-holding pause, may be adjusted independently to maximize control over the deposition site and level.

# Dynamical analysis of an HIV infection model including quiescent cells and immune response

Nali Ibrahim<sup>\*1</sup>, Attila Denes<sup>2</sup> and Xueyong Zhou<sup>3</sup>

<sup>1</sup> University of Szeged, Bolyai insitute, Hungary

<sup>2</sup> University of szeged, Bolyai insitute, Hungary

<sup>3</sup> Xinyang Normal University, China

Email : ibrahim.nali@usmba.ac.ma

## Abstract

This paper presents a comprehensive analysis of an HIV infection model that incorporates quiescent cells and immune response dynamics within the host. The model, represented by a system of ordinary differential equations, captures the intricate interplay between the hosts immune response and the viral infection. The study investigates the fundamental properties of the model, including equilibrium analysis, the computation of the basic reproduction number  $\mathcal{R}_0$ , stability analysis, bifurcation phenomena, numerical simulations, and sensitivity analysis. An infection equilibrium, which reflects the persistence of the infection, and a disease-free equilibrium, which represents the possibility of disease control, are both revealed by the analysis. By applying matrix-theoretical methods, stability analysis confirmed that the disease-free equilibrium is both locally and globally stable for  $\mathcal{R}_0 < \infty$ . The research also reveals a transcritical forward-type bifurcation at  $\mathcal{R}_0 = \infty$ , which denotes a critical threshold that affects the behavior of the system. The temporal dynamics of the model are investigated through numerical simulations, and sensitivity analysis determines the most important variables by examining the effects of parameter changes on the systems behavior.

## References

- [1] BOCHAROV, G., CHERESHNEV, V., GAINOVA, I., et al. Human immunodeficiency virus infection: from biological observations to mechanistic mathematical modelling. *Mathematical Modelling of Natural Phenomena* 7.5 (2012): 78104.
- [2] Dorratohtaj, Nargesalsadat, et al. Multi-scale immunoepidemiological modeling of within-host and between-host HIV dynamics: systematic review of mathematical models. *PeerJ* 5 (2017): e3877.
- [3] Perelson, Alan S., and Ruy M. Ribeiro. Modeling the within-host dynamics of HIV infection. *BMC biology* 11 (2013): 110.

# Fractional ordered model for cell level viral transmission dynamics with adaptive immunity

Getachew Teshome Tilahun\*<sup>1</sup>

<sup>1</sup> Haramaya University, Department of Mathematics

Email : gmgech@gmail.com

## Abstract

In this paper we proposed fractional ordered model of cell level pathogen transmission dynamics by considering adaptive immunity. Infected and latently infected cells are considered in the model. The model assumed as infection occur from virus to health cell and from infected cell to health cells. The qualitative behavior of the model is analyzed. From the qualitative analysis basic threshold parameters that affect clearance and maintenance of infection are determined. Effect memory as well as some parameters on the expansion of the infection is done in numerical simulations.

**Keywords:** Fractional Order; Modeling; Cell level; Adaptive immunity

## The SEIR model With Unreported Cases and Markovian Switching

H. J. Alsakaji<sup>1</sup>, Y. A. El-Khatib<sup>\*1</sup> and F.A. Rihan<sup>1</sup>

<sup>1</sup>Department of Mathematical Sciences, College of Science, United Arab Emirates University, Al Ain, 15551, UAE

Email : [youssef\\_elkhatib@uaeu.ac.ae](mailto:youssef_elkhatib@uaeu.ac.ae)

### Abstract

The study of infectious diseases in humans has become increasingly important in public health. This paper extends the SEIR model to include unreported cases (U), environmental white noise, and the impact of an exogenous event. Environment variation is used to conduct dynamic analysis. We examine the process of switching from one regime to another at random. The ergodicity and stationary distribution criteria are discussed. Using a Lyapunov function, we write down some sufficient conditions for disease extinction. With different intensities of stochastic noises, we calculate the threshold of extinction for the stochastic epidemic system. In order to control the spread of disease, stochastic noise plays an important role. A numerical simulation and a fit to real data have shown that the model and theoretical results are valid.



# Mathematical modeling of the effect of quarantine rate on controlling the infection of COVID19 in the population of Saudi Arabia

Ahuod S. Alsheri<sup>\*1</sup>, Aeshah A. Alraeza<sup>2</sup> and Mona R. Afia<sup>1</sup>

<sup>1</sup> Department of Mathematics, University of Bisha, 8154, Alnamas 67398-5644, Saudi Arabia.

<sup>2</sup> Department of Mathematics, King Khalid University, Abha 62529, Saudi Arabia

Email : oshokry@ub.edu.sa

## Abstract

With the development of communications and transportation worldwide, the challenge of controlling epidemiological diseases becomes higher. The COVID19 has put all nations in a lethal confront with a severe disease that needed serious and painful actions. The sooner the actions, the less destructive the impact. In this paper, we incorporate what we believe is crucial but applicable to control the spread of COVID19 in the populations, that is, quarantine. We keep the model as simple as SI Kermack-McKendrick model with an additional compartment of quarantined patients. We established the systems basic properties and studied the stability of the disease-free equilibrium and its relation to the basic reproduction number  $\mathcal{R}$ , in which we calculated its formula. The focus of our study is to measure the effect of quarantine rate on controlling the spread of COVID19. We use the data collected from the Ministry of Health in Saudi Arabia. We studied three different values of the quarantine rate where newly infectious patients are detected and isolated within 14, 7, and 5 days. The simulations show a significant effect of the quarantine where COVID19 can be fully controlled if the newly infected patient enters the quarantine within five days. These results were proposed to the Public Health Authority in Saudi Arabia and approved by the Ministry of Health in which they applied promptly.

# Optimal control for cancer-tumor-immune systems using reinforcement learning

R. RAKKIYAPPAN<sup>\*1</sup>

<sup>1</sup> Department of Mathematics, Bharathiar University, Coimbatore641046, India.

Email : rakkigru@gmail.com

## Abstract

In this work, a class of cancer-tumor-immune systems with drug dosage acting as control is described as [1, 2]

$$\begin{aligned}\dot{T} &= \mu_c TF(T) - \gamma TI - \kappa_T Tu \\ \dot{I} &= \mu_I (T - \beta T^2) I - \delta I + \alpha + \kappa_I Iv\end{aligned}$$

In the above set of equations, the values T represents the tumor volume and I denotes the immune competent cell densities. The ability of the immune cells to eliminate the tumor cells are denoted through the tumor-cell-elimination rate  $\gamma \in \mathbb{R}$ . The positive constants  $\alpha, \delta, \beta, \mu_I$  denotes the T-cells generation rate, the natural death rate of immune cells and calibrations these interactions and the way cancer cells influence the stimulation of immune cells, respectively. The control input u determined though drug dosage eliminates the tumor cells. is utilized to denote the booster drug for the immune system. The positive coefficients  $\kappa_T$  and  $\kappa_I$  represent the control capability of the control activity of drug dosage and immune system booster drug, respectively. An optimal control for cancer-tumor-immune systems is discussed using reinforcement learning

## References

- [1] Ledzewicz, U., Naghnaeian, M., & Schattler, H. (2010, December). Bifurcation of singular arcs in an optimal control problem for cancer immune system interactions under treatment. In 49th IEEE Conference on Decision and Control (CDC) (pp. 7039-7044). IEEE.
- [2] Stepanova, N. V. (1979). Course of the immune reaction during the development of a malignant tumour. *Bio-physics*, 24(5), 917-923

# Hepatitis B epidemic model: analysis and control

Fathalla A. Rihan<sup>1</sup> and Tahir Khan <sup>\*1</sup>

<sup>1</sup>UAE University, Al-Ain, P.O. Box 15551, United Arab Emirates

Email : tahir k@uaeu.ac.ae

## Abstract

The infection of hepatitis B produces inflammation of the liver and can cause chronic and acute diseases, while it is life-threatening and a major health problem around the globe. The control of this infection is a challenging task because of many reasons such as proper medication, variation of human behavior, vaccination, and particularly the existence of a large number of carriers, however, understanding the temporal dynamics of the infection helps to present appropriate control mechanisms. Thus, a complex dynamical system is needed to find the stability conditions and propose various intervention strategies to forecast the control of the contagious infection of hepatitis B. We develop a model to investigate the temporal dynamics and suggest a control mechanism for the infection of hepatitis B. The well-posedness will be shown to prove the feasibility of the proposed epidemic problem and find the threshold parameter to analyze the model equilibria and its stability. We also discuss the sensitive analysis of the reproductive number to quantify the parameters that are more sensitive to disease spreading and control. Based on the model dynamics and sensitivity, we investigate effective procedures to minimize the infection of hepatitis B and develop algorithms to verify the theoretical work with the help of numerical experiments.

# Comprehensive cost-effectiveness analysis of a new compartmental model for bacterial meningitis considering the influence of the media

Joshua Kiddy K. Asamoah<sup>\*1</sup>

<sup>1</sup> Department of Mathematics, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana

Email : [jkkasamoah@knust.edu.gh](mailto:jkkasamoah@knust.edu.gh)

## Abstract

Mathematical epidemiology has paid some attention to the study of bacterial meningitis because of the severity of the disease and the way it spreads through a population. In this paper, we formulated a new compartmental model for meningitis to study the impact of media on the reduction of the severity of the disease in Ghana. We obtain the control reproduction number and the herd immunity rate in the presence of vaccination. We noticed that the meningitis death rate can be control with an increase in publicity of the dynamics of the spread of the disease and the ability to seek immediate medical treatment. We conduct a sensitivity analysis using Latin hypercube sampling. It is noticed that the rate of transmission and that of, vaccination and the media have a negative nonlinear correlation. The effects of media, vaccination, and treatment as a function of time are also investigated using an optimal control model. We display the efficiencies of the controls and demonstrate the mean and incremental cost-effectiveness ratios. The cost of controlling meningitis in the presence of media, vaccination, and treatment is discussed, and we hope that this study will help to further educate the public on this topic.

**Keywords:** Control Reproduction Number, Media, Vaccination, Treatment, Cost-effectiveness analysis

# Dynamics of HIV-COVID-19 Co-Infection: A Fractional Order Model and Analysis using Laplace-Adomian Decomposition Method

Isa Abdullahi Baba<sup>\*1,2</sup>, and Fathalla A. Rihan<sup>2,3</sup>

<sup>1</sup> Department of Mathematical Sciences, Bayero University, Kano, Nigeria

<sup>2</sup> Department of Mathematical Sciences, College of Science, UAE University, Al Ain 15551, UAE

<sup>3</sup> Department of Mathematics, Faculty of Science, Helwan University, Cairo 11795, Egypt

Email : iababa.mth@buk.edu.ng

## Abstract

The co-infection of HIV and COVID-19 presents a significant health challenge with potentially severe consequences. Understanding the dynamics of HIV-COVID-19 co-infection is crucial for developing effective control measures and optimizing healthcare strategies. In this study, we develop a mathematical model based on Caputo fractional order differential equations to capture the complex dynamics of co-infection. The model incorporates the transmission dynamics of HIV and COVID-19, the immune response of the host, and the effects of treatment interventions. To solve the fractional order model, we employ the Laplace Adomian decomposition method, a powerful technique for obtaining approximate solutions of fractional order differential equations. By simulating the interactions between these factors, we gain insights into the spread of coinfection and identify critical factors contributing to its progression. Additionally, we analyze the convergence properties of the series solution obtained using the Laplace Adomian decomposition method. Our findings will inform the development of effective control measures and assist policymakers, healthcare providers, and public health authorities in mitigating the spread and impact of HIV-COVID-19 co-infection.

**Keywords:** COVID-19; HIV; Co-infection; Laplace Adomian decomposition method; series solution; convergence.

# Traveling wave solutions on synergistic co-infection model in crop diseases

AMERICO MATUSSE\*<sup>1</sup>

<sup>1</sup> University of Pretoria, Lynnwood Rd, Hatfield, Pretoria, 0002, South Africa

Email : ajmatusse@gmail.com

## Abstract

Plants and crops are repeatedly defied by pathogens during their lifespan, causing diseases with a negative impact on the host fitness. In particular, pathogens have caused drastic yield reductions in several important crops. For instance, FAOSTAT database indicates that annually, an yield losse of 20%-40% is due to pests. The majority of plants and crops can be infected by two or more unrelated viruses, leading to more severe symptoms, such as in the case of Maize Lethal Necrosis Disease (MLND) as a synergistic co-infection of Maize Chlorotic Mottle Virus (MCMV) and a potyvirus. See for instance [1, 2]. In this talk, we develop a general spatial explicit synergistic coinfection mathematical model for vector and plant of two virus. The model takes into account the infection through virus only. We then discuss theoretically and numerically the existence of traveling waves solutions. Finally, we discuss possible invasion control through numerical simulation.

## References

- [1] Chapwanya, M., Matusse, A., Dumont, Y., On synergistic co-infection in crop diseases. The case of the Maize Lethal Necrosis Disease, *Applied Mathematical Modelling*, 90, 912-942, 2021.
- [2] Hilker, F. M., Allen, L. J., Bokil, V. A., Briggs, C. J., Feng, Z., Garrett, K. A., ... Cunniffe, N. J., Modeling virus coinfection to inform management of maize lethal necrosis in Kenya, *Phytopathology*, 107(10), 1095-1108, 2017.

# Global analysis of an SEI plant disease model with saturation incidence rate

Debasis Mukherjee<sup>1</sup>

<sup>1</sup>Department of Mathematics Vivekananda College Thakurpukur Kolkata-700063 India

Email : mukherjee1961@gmail.com

## Abstract

This paper studies an SEI epidemic model for plant disease with saturation incidence rate. Positivity and boundedness of solutions are examined to justify the model biologically. Local stability condition of boundary as well as endemic (positive) equilibrium point is derived. Persistence criterion is developed. The existence of a Hopf bifurcation is shown by considering the intrinsic growth rate as well as disease transmission rate as a bifurcation parameter. Applying Bendixsons criterion for high-dimensional ordinary differential equations, we prove that the endemic equilibrium point is globally stable under certain conditions. The dynamical behaviour of the model is highly sensitive to fluctuations in the disease transmission rate. Finally, numerical simulations are carried out to validate the results obtained.

# A novel investigation of the tuberculosis disease via fractional differential operator

SHYAMSUNDER\*<sup>1</sup>

<sup>1</sup>Department of Mathematics, Malviya National Institute of Technology Jaipur, India

Email : skumawatmath@gmail.com

## Abstract

In this study, a computational solution to the fractional tuberculosis disease is obtained with the help of the Generalized Eulers Method [1, 2]. This article considers two treatment strategies: protective therapy for latent populations and main treatment for infected populations. The compartmental structure of the six-dimensional model includes the susceptible, latent, infected, recovered, and treatment classes. Additionally, we studied the stability of the equilibrium point. To examine the nature of multidimensional differential equations of fractional order that arise in biological illness, the findings produced utilizing the method under consideration are more accurate and straightforward. Additionally, we studied the stability of the equilibrium point. The fractional model has been graphically simulated using MATLAB22.

## References

- [1] Kilbas A.A., Srivastav H.M., Trujillo J.J., Theory and applications of fractional differential equations, Elsevier, 204, 2006.
- [2] Odibat Z.M., Momani S., An algorithm for the numerical solution of differential equations of fractional order, Journal of Applied Mathematics & Informatics, 26(1), 1527, 2008.



# Mathematical Analysis and Parameter Estimation of a Two-Patch Zika Model

Kifah Almaqrashi\*<sup>1</sup>

<sup>1</sup>Sultan Qaboos University

Email : Kifahhsm@gmail.com

## Abstract

In this work, we developed a multi-patch model for the spread of Zika virus infection taking into account direct and indirect transmissions along with vertical transmission. The model was analyzed to gain some insights into the spread of the disease. The model was fitted to a data set collected from two neighboring countries Brazil and Colombia in order to estimate some of its parameters and use it for calculating the reproduction number and for the purpose of sensitivity analysis. Our results show that the reproduction number is less than one in both countries, which indicates that the disease will die out. Also, our results show that direct transmission is the most important route for the spread of the disease, and hence it has to gain more focus in any controlling strategy.

## Impact of fear of a fractional-order plant disease model with herbivore attack

CHANDAN MAJI\*<sup>1</sup>

<sup>1</sup> Jagannath Kishore College

Email : chandanmaji.ju@gmail.com

### Abstract

This work deals with a fractional-order mathematical model for plants subject to a disease and harvested by herbivores. The existence, uniqueness, non-negativity and boundedness of the considered model has been analyzed. A detailed analysis of different equilibria and their stability has been discussed. Global stability of the interior equilibrium point is investigated. It has been shown that fear factor and fractional order have an important role in the dynamics of system stability. Finally, some numerical simulations are performed to validate our findings.

# Optimal heat transport induced by magnetic nanoparticle delivery in vascularised tumours

Tahani Al Sariri\*<sup>1</sup>

<sup>1</sup> Sultan Qaboos University

Email : alsariri@squ.edu.om

## Abstract

We describe a novel mathematical model for blood flow, delivery of nanoparticles, and heat transport in vascularised tumour tissue. The model, which is derived via the asymptotic homogenisation technique, provides a link between the macroscale behavior of the system and its underlying, tortuous micro-structure, as parametrised in Penta and Ambrosi (2015) [1]. It consists of a double Darcys law, coupled with a double advectiondiffusionreaction system describing heat transport, and an advectiondiffusionreaction equation for transport and adhesion of particles. Particles are assumed sufficiently large and do not extravasate to the tumour interstitial space but blood and heat can be exchanged between the two compartments. Numerical simulations of the model are performed using a finite element method to investigate cancer hyperthermia induced by the application of magnetic field applied to injected iron oxide nanoparticles. Since tumour microvasculature is more tortuous than that of healthy tissue and thus suboptimal in terms of fluid and drug transport, we study the influence of the vessels geometry on tumour temperature. Effective and safe hyperthermia treatment requires tumour temperature within certain target range, generally estimated between  $42^{\circ}C$  and  $46^{\circ}C$ , for a certain target duration, typically 0.5h to 2h. As temperature is difficult to measure in situ, we use our model to determine the ranges of tortuosity of the microvessels, magnetic intensity, injection time, wall shear stress rate, and concentration of nanoparticles required to achieve given target conditions.

## References

- [1] Penta, R., Ambrosi, D., 2015. The role of the microvascular tortuosity in tumor transport phenomena. *J. Theoret. Biol.* 364, 8097.

# Traveling wave speed and profile of a go or grow glioblastoma multiforme model

Aisha TURSINKOZHA<sup>1</sup>, Ardak KASHKYNBAYEV\*<sup>1</sup>, Bibinur SHUPEYEVA<sup>1</sup>, Erica M. RUTTER<sup>2</sup> and Yang KUANG<sup>3</sup>

<sup>1</sup>Department of Mathematics, Nazarbayev University, Astana, Kazakhstan

<sup>2</sup>Department of Applied Mathematics, University of California, Merced, CA, USA

<sup>3</sup>School of Mathematical and Statistical Sciences, Arizona State University, Tempe, AZ, USA

Email: ardak.kashkynbayev@nu.edu.kz

## Abstract

Glioblastoma multiforme (GBM) is a fast-growing and deadly brain tumor due to its ability to aggressively invade the nearby brain tissue. A host of mathematical models in the form of reaction-diffusion equations have been formulated and studied in order to assist clinical assessment of GBM growth and its treatment prediction. To better understand the speed of GBM growth and form, we propose a two-population reaction-diffusion GBM model based on the go or grow hypothesis. Our model is validated by in vitro data and assumes that tumor cells are more likely to leave and search for better locations when resources are more limited at their current positions. Our findings indicate that the tumor progresses slower than the simpler Fisher model, which is known to overestimate GBM progression. Moreover, we obtain accurate estimations of the traveling wave solution profiles under several plausible GBM cell switching scenarios by applying the approximation method introduced by Canosa.

## References

- [1] Tursynkozha, A., Kashkynbayev, A., Shupeyeva, B., Rutter, E.M., Kuang, Y, Traveling wave speed and profile of a go or grow glioblastoma multiforme model, *Communications in Nonlinear Science and Numerical Simulation*, **118** (2023), 107008

## In-depth epidemiology and modeling of the future trends of communicable and non-communicable diseases in Kazakhstan using aggregated big data from the Unified National Electronic Healthcare System

Abduzhappar GAIPOV\*<sup>1</sup>, Gulnur ZHAKHINA<sup>1</sup>, Sauran YERDESSOV<sup>1</sup>, Yesbolat SAKKO<sup>1</sup>, Temirgali AIMYSHEV<sup>1</sup>, Kamilla MUSSINA<sup>1</sup>, Dmitriy SYSSOYEV<sup>1</sup>, Iliyar ARUPZHANOV<sup>2</sup>, Amin ZOLLANVARI<sup>2</sup> and Ardak KASHKYNBAYEV<sup>3</sup>

<sup>1</sup> Department of Medicine, School of Medicine, Nazarbayev University, Astana, Kazakhstan

<sup>2</sup> Department of Electrical and Computer Engineering, School of Engineering and Digital Sciences, Nazarbayev University, Astana, Kazakhstan

<sup>3</sup> Department of Mathematics, School of Sciences and Humanities, Nazarbayev University, Astana, Kazakhstan

Email : aabduzhappar.gaipov@nu.edu.kz

### Abstract

There is an emerging need to develop a comprehensive epidemiological and indepth analysis of communicable and non-communicable diseases in Kazakhstan. Monitoring electronic health systems (UNEHS) and digitalization of healthcare, disease registries, including most of the emerging diseases that Kazakhstan has implemented. This study has had ultimately aims for the development of a research center for disease control and prevention, using modern epidemiological analytic approaches, advanced biostatistical evaluations, mathematical modeling and machine learning to support with appropriate epidemiologic and methodologic techniques to working on large-scale administrative healthcare data for public health surveillance of infectious diseases. features, particularly, geographical, seasonal and other patterns of communicable and non-communicable diseases using UNEHS. Along with statistical tools, we used mathematical modeling and machine learning algorithms which have provided a better and more comprehensive understanding of the infection transmission dynamics and forecasting the trend of the diseases. Several peer reviewed papers have been published as well as still ongoing many research projects using data from UNEHS and different methodological approaches such as advanced epidemiology, mathematical modeling, and machine learning.

**Funding:** This study was supported by grants from the Nazarbayev University Faculty Development Aggregation and utilization of the large-scale administrative health data in Kazakhstan for population health research and surveillance), Nazarbayev University Faculty Development Research Grant Program modeling of the 10-year trends of cardiovascular diseases and their complications in Kazakhstan using aggregated big data from the united National Electronic Healthcare System) and grant from Ministry

**Keywords:** Big Data in Healthcare, Epidemiology, Machine Learning, Mathematical models

## **References**

- [1] Gusmanov, A., Zhakhina, G., Yerdessov, S., Sakko, Y., Mussina, K., Alimbayev, A., Syssoyev, D., Sarria-Santamera, A., & Gaipov, A. , Review of the research databases on population-based for epidemiological research and Real-World Evidence, *International Journal of Medical Informatics*, 170 (2022), 104950
- [2] Mussina, K., Kadyrov, S., Kashkynbayev, A., Yerdessov, S., Zhakhina, G., Sakko, Y., Zollanvari, A., & Gaipov, A. Prevalence of HIV in Kazakhstan 2010-2020 and Its Forecasting for the Next 10
- [3] Alimbayev, A., Zhakhina, G., Gusmanov, A., Sakko, Y., Yerdessov, S., Arupzhanov, I., Kashkynbayev, A., Zollanvari, A., & Gaipov, A., Predicting 1-year mortality of patients with diabetes *Reports*, 13(1) (2023)
- [4] Zhakhina, G., Gaipov, A., Salustri, A., Gusmanov, A., Sakko, Y., Yerdessov, S., Bekbossynova, M., Abbay, A., Sarria-Santamera, A., & Akbilgic, O., Incidence, mortality and disability-adjusted life

## Multi-stability analysis of fractional-order quaternion-valued neural networks with time delay

KATHIRESAN S<sup>\*1</sup>, ARDAK KASHKYNBAYEV<sup>1</sup>, K. JANANI and R. RAKKIYAPPAN<sup>2</sup>

<sup>1</sup>DEPARTMENT OF MATHEMATICS, NAZARBAYEV UNIVERSITY, NUR-SULTAN  
010000, KAZAKHSTAN

<sup>2</sup> DEPARTMENT OF MATHEMATICS, BHARATHIAR UNIVERSITY, COIMBATORE - 641  
046, TAMILNADU, INDIA

Email : kathiresan.sivakumar@nu.edu.kz

### Abstract

This paper addresses the problem of multi-stability analysis for fractional-order quaternion-valued neural networks (QVNNs) with time delay. Based on the geometrical properties of activation functions and intermediate value theorem, some conditions are derived for the existence of at least  $(2\mathcal{K}_p^R + 1)^n$ ,  $(2\mathcal{K}_p^I + 1)^n$ ,  $(2\mathcal{K}_p^J + 1)^n$ ,  $(2\mathcal{K}_p^K + 1)^n$  equilibrium points, in which  $[(\mathcal{K}_p^R + 1)^n, (\mathcal{K}_p^I + 1)^n, (\mathcal{K}_p^J + 1)^n, (\mathcal{K}_p^K + 1)^n]$  of them are uniformly stable while the other equilibrium points become unstable. Thus the developed results show that the QVNNs can have more generalized properties than the real-valued neural networks (RVNNs) or complex-valued neural networks (CVNNs). Finally, two simulation results are given to illustrate the effectiveness and validity of our obtained theoretical results.

### References

- [1] F. Zhang, Z. Zeng, Multistability and instability analysis of recurrent neural networks with time-varying delays, *Neural Netw.*, 97 (2018), 116126. doi: <https://doi.org/10.1016/j.neunet.2017.09.013>.
- [2] Z. Zeng, T. Huang, W X. Zheng, Multistability of recurrent neural networks with time-varying delays and the piecewise linear activation function, *IEEE Trans. Neural Netw. Learn. Syst.* 21 (2010), 13711377. doi: 10.1109/TNN.2010.2054106.

# A New Investigation on the Impact of Vaccination on the Transmission of Monkeypox

M. Manivel\*<sup>1</sup> and A. Venkatesh<sup>1</sup>

<sup>1</sup> Department of Mathematics, AVVM Sri Pushpam College, (Affiliated to Bharathidasan University), Poondi, Thanjavur, Tamilnadu, India-613 503.

Email : manivelmani718@gmail.com

## Abstract

The primary aim of this study is to construct a fuzzy fractional mathematical model to investigate the dynamics of monkeypox viral transmission. This dynamical model of monkeypox vaccinations humans is developed using Caputos fuzzy fractional differential equation. Our technique applies to the fuzzy system of fractional ordinary differential equations, and our findings are correct. The impact of the fractional operator on the different compartments of the model has been observed through numerical simulation. The findings demonstrate that maintaining a vaccinated human population mitigates disease transmission. We conclude with short comments and numerical simulations to demonstrate the results.

**Keywords:** monkeypox virus, fractional derivative, fuzzy fractional differential equation, Caputos differential equation.



## Finite-time/fixed-time synchronization of memristive shunting inhibitory cellular neural networks via sliding mode control

Madina Otkel<sup>\*1</sup>, Soundararajan Ganesan<sup>1</sup>, Rakkiyappan Rajan<sup>2</sup> and Ardak Kashkynbayev<sup>1</sup>

<sup>1</sup>Nazarbayev University, Astana, Kazakhstan

<sup>2</sup>Bharathiar University, Tamil Nadu, India

Email : amadina.otkel@nu.edu.kz

### Abstract

This abstract proposes a novel time-dependent gain parameter-based sliding mode controller (SMC) to realize the finite-time/fixed-time synchronization of memristive shunting inhibitory memristive neural networks (Mem-SICNNs) having time-varying delays. In this regard, a new terminal sliding mode surface is designed and its reachability is analysed. According to synchronization error analysis, the stability property of the desired error system is reached within finite-time/fixed-time range by proposing a unique time-dependent gain parameter-based SMC and choosing the appropriate Lyapunov functionals. Finally, a numerical example is approached by software simulation and manual calculation to estimate the settling-time of the finite-time/fixed-time synchronization criteria of the proposed Mem-SICNNs model.

**Funding:** The authors were supported by the Science Committee of the Ministry of Science and Higher Education of the Republic of Kazakhstan (Grant No.: AP13068282 Synchronization and anti-synchronization analysis of discrete-time neural networks with delays) and by Nazarbayev University under Collaborative Research Program Grant No: 11022021CRP1509.

**Keywords:** finite-time/fixed-time synchronization; memristor; shunting inhibitory cellular neural networks; sliding mode control.

## Structured dynamics of the cell-cycle at multiple scales

Arran Hodgkinson<sup>1</sup>, Aisha Tursynkozha<sup>\*2</sup> and Dumitru Trucu<sup>3</sup>

<sup>1</sup>Quantitative Biology and Medicine, Living Systems Institute, University of Exeter, Exeter, United Kingdom

<sup>2</sup>Department of Mathematics, Nazarbayev University, Astana, Kazakhstan

<sup>3</sup>Division of Mathematics, University of Dundee, Dundee, United Kingdom

Email: aisha.tursynkozha@nu.edu.kz

### Abstract

The eukaryotic cell cycle consists of four distinct phases ( $G_1$ ,  $S$ ,  $G_2$ , and  $M$ ), and it ensures the protection of the cell from DNA-damage and becoming a cancerous cell. DNA-damage activates the regulatory protein p53, which arrests the cell cycle to allow for the repair of this damage, while, p53 is a single inactivated and frequently mutated protein in cancer. As such, this procedure takes place at the cellular level; however, its impact is observed at the level of the entire organism's cell population. In this context, we introduce two models of cell cycle progression: The first model focuses on the cell-scale process of cell cycle progression and the temporal biochemical processes, driven by cyclins, which govern the transition from one phase to another. On the other hand, the second model addresses the broader aspect of cell-cycle progression across a population of cells and its arrest caused by DNA-damage and activation of the p53 gene.

# Asymptotic analysis of Alzheimers disease in thin heterogeneous domains

ACHILLE LANDRI POKAM KAKEU<sup>\*1</sup>

<sup>1</sup> Department of Mathematics and Computer Science, University of Dschang, P.O. Box 67, Dschang, Cameroon

Email : pokakeu@yahoo.fr

## Abstract

We carry out in thin layer the homogenization of a set of Smoluchowskis discrete diffusion-coagulation equations describing the evolution density of diffusion particles that are prone to coagulate in pairs. Alzheimers disease (AD), the most common cause of dementia, destroys cognitive skills and memory as a progressive neurodegenerative disease. The aggregation and the diffusion of  $\beta$ -amyloide peptide ( $A\beta$ ) are responsible to the development of Alzheimers disease at different scales and the thin heterogeneous domains give rise to these phenomena in lower space dimensions. More precisely, assuming that the thin heterogeneous layer is made of microstructures that are distributed inside in a deterministic way including as special cases the periodic and the almost periodic distributions, we make use of the concept of algebras with mean value to state the main compactness results and we also make use of a new approach of the sigma-convergence method that is suitable for the study of such phenomena occurring in thin heterogeneous media. This is made through a systematic study of the sigma-convergence method for thin heterogeneous domains.

## References

- [1] Yves Achdou, Bruno Franchi, Norina Marcello, and Maria Carla Tesi. A qualitative model for aggregation and diffusion of beta-amyloid in alzheimers disease. *Journal of mathematical biology*, 67(6):1369- 1392, 2013.
- [2] Pokam Kakeu, A. L. and Woukeng, J. L. (2020), Homogenization of nonlinear parabolic equations with hysteresis, *ZAMM-Journal of Applied Mathematics and Mechanics/Zeitschrift fur Angewandte Mathematik und Mechanik* p. e201900323.
- [3] A. L. Pokam Kakeu, J.L. Woukeng, Well-Posedness and Long-Time Behaviour for a Nonlinear Parabolic Equation With Hysteresis, *Commun. Math. Anal.* 23 (2020) 3862.

# Control Analysis of Nilaparvata Lugens with Wolbachia using Sterile Insect Techniques

Rajivganthi Chinnathambi\*<sup>1</sup>

<sup>1</sup>Vellore Institute of Technology, Chennai Campus

Email : rajivganthi.c@vit.ac.in

## Abstract

Modeling and controlling Nilaparvata Lugens (Lugens) populations in paddy fields have been highly beneficial to the environment and farmers alike. As part of this study, we describe a sex-structured wild Lugens model and male Lugens infected with w-Stri (Wolbachia). A laboratory test has shown that w-Stri type Wolbachia is able to control wild Lugens biologically. When male Lugens infected with w-Stri mate with wild female Lugens, they produce higher levels of cytoplasmic incompatibility (CI). We construct an optimal control problem by using the continuous releasing rate of male Lugens infected with w-Stri as a time-dependent control parameter. We examine the optimality in terms of its necessary and sufficient conditions. Furthermore, we discuss the sufficient conditions to reach the elimination of wild Lugens through periodic impulsive releases of male Lugens infected with wStri. Numerical simulations are performed to validate the theoretical conclusions.

# Cinematographic Method for Mathematical Modeling of Heat Transfer and Exchange Processes in Geometrically Complicated Bodies

Abdulla Azamov\*<sup>1</sup>

<sup>1</sup> Institute of Mathematics Uzbekistan Academy of Sciences

Email : [abdulla.azamov@gmail.com](mailto:abdulla.azamov@gmail.com)

## Abstract

Heat distribution, transfer and exchange processes are widely spread as in nature so in technics. If a region where such a process is to be considered has simple geometric features as prisms, cylinders, balls, classic models mathematical models are effective. In cases when the process occurs in complicated bodies it is preferable an approach of discretization simulating cinematographic motion. This approach allows to calculate heat distribution in metallic grid of a rotating air preheater of energy plants and heat exchange between tube and gas flowing in it when the tube has arbitrary complicated configuration. The main point of the method is a reduction of the problem to linear systems of discrete equations while the most difficulty arises in solving the inverse problem in order to identify the model. It will be discussed questions of developing the method taking into account delay in time and control parameters and game problems.

## References

- [1] A. A. Azamov, M. A. Bekimov. Simplified model of the heat exchange process in rotary regenerative air preheaters. *Ural Mathematical Journal*, Vol. 2 (2016), No. 2, P. 12-19.
- [2] Bekimov M. A., F. A. Rihan. A discrete mathematical model for a heat transfer process in rotating regenerative air preheater. *Differential equations and dynamical systems. USUZCAMP 2017. Springer proceedings in mathematics statistics*, 2018, vol. 268. Springer, Cham. pp. 55-61.

# A continuum space is the infinitely great

Qing Li\*<sup>1</sup>

<sup>1</sup> ShiJiaZhuang Traditional Chinese Medical Hospital, ShiJiaZhuang, HeBei province, China

Email : liqingliyang@126.com

## Abstract

An infinitely small quantity is defined as a one-dimensional quantity of finite length but with sizes of space, while an infinitely great quantity is reached by the superposition or accumulation of infinitely many finite quantities, by the way of the change in direction. The change in direction indicates that there is a jump from a finite quantity to infinitely many finite quantities (infinitely great). The form of the manifestation of the infinitely great is one quantitative continuum that cannot be operated by any algorithms and all parts of space we see is this one quantitative continuum. Any value are this single quantitative continuum that indicates the infinitely great and compresses any quantities outside of it to nothing. As a result, the infinite exact value of a circumferential length ( $\pi$ ) has been obtained here.

**Keywords:** infinitely great, change in direction, one quantitative continuum

## References

- [1] Qing Li .A geomerty consisting of singularities containing only integers. (preprint Research Square: DOI: 10.21203/rs.3.rs-219046/v1)
- [2] Wiles A, Modular elliptic curves and Fermats Last Theorem., *Annals of Mathematics*, vol.141, pp. 443-552, 1995.
- [3] Jeremy Kahn and Vladimir Markovic, Immersing almost geodesic surfaces in a closed hyperbolic three manifold, *Annals of Mathematics*, vol.175, pp.1127-1190, 2012.
- [4] Christoph Bhm and Burkhard Wilking, Manifolds with positive curvature operators are space forms, *Annals of Mathematics* ,vol.167, pp.1079-1097 ,2008.
- [5] Jeff Cheeger and Aaron Naber, Regularity of Einstein manifolds and the codimension 4 conjecture, *Annals of Mathematics* ,vol.182, pp.1093-1165 ,2015.

## Degree Reduction of Bezier Curves

Abdallah Rababah<sup>1</sup>

<sup>1</sup>Department of Mathematical Sciences, College of Science, United Arab Emirates University, Al Ain 15551, UAE

Email : rababah@uaeu.ac.ae

### Abstract

There are substantial methods of degree reduction in the literature. Existing methods share some common limitations, such as lack of geometric continuity, complex computations, and one-degree reduction at a time. In this talk, an approximate geometric multi-degree reduction algorithm of Bezier curves is proposed. Geometric continuity conditions are applied in the degree reduction process to preserve the boundary control points. The general equation for high-order (G2 and above) multi-degree reduction algorithms is nonlinear, and the solutions of these nonlinear systems are quite expensive. In this paper,  $G_1$ -continuity conditions are imposed besides the  $G_0$ -continuity conditions. While some existing methods only achieve the multi-degree reduction by repeating the one-degree reduction method recursively, the proposed method achieves multi-degree reduction at once. The distance between the original curve and the degree-reduced curve is measured with the 1-infinite norm. Numerical examples and figures are presented to state the adequacy of the algorithm. The proposed method not only outperforms the existing methods of degree reduction but also guarantee geometric continuity conditions at the boundary points, which is very important in CAD and geometric modeling.

## Role of the NFAT-signaling network in controlling allergic skin reactions

Muhammad Azeem<sup>1</sup>, Andreas Kerstan<sup>1</sup> and Khalid Muhammad\*<sup>2</sup>

<sup>1</sup> Department of Dermatology, University Hospital Wrzburg, Germany

<sup>2</sup>UAE University, Al Ain, United Arab Emirates

Email : k.muhammad@uaeu.ac.ae

### Abstract

Allergic contact dermatitis (ACD) is a type of human skin allergic reactions caused by multiple contact allergens. It accounts for at least 20% which stresses on the need of basic and clinical research on it. So far, a plethora of data supports a major role of CD4+ and CD8+ T lymphocytes in the progression and maintenance of disease. However, in-depth understanding about the molecular mechanisms of its pathophysiology is still limited. The Ca<sup>++</sup>-Calcineurin-NFAT-signalling network is an essential mediator of T cell receptor signaling. Out of four NFATc members, NFATc1 plays a prominent role in T cell activation. Therefore, we investigated its role in contact hypersensitivity (CHS) which is an established animal model to study human ACD. The 2,4,6- trinitrochlorobenzene (TNCB)-induced CHS response was diminished in *Nfatc1<sup>fl/fl</sup>Cd4-cre* mice (*Nfatc1<sup>-/-</sup>*) as compared to wild-type (WT) mice. The reduced CHS response was linked to a lower percentage of interleukin (IL)17-producing CD8+ T (Tc17) cells in inflamed skin and draining lymph nodes (dLN). In vitro Tc17 polarization assays displayed that *Nfatc1<sup>-/-</sup>* CD8+ T cells have a reduced ability to polarize into Tc17 cells. In the absence of NFATc1, CD8+ T cells favor the differentiation of Interferon (IFN)- $\gamma$  secreting CD8+ T (Tc1) cells, whereas in its presence they develop into Tc17 cells. Moreover, the CHS response was rescued in nave *Nfatc1<sup>-/-</sup>* mice by adoptive transfer of TNCB-sensitized WT CD8+ T cells. Finally, we detected the infiltration of Tc17 cells in human ACD skin samples by immunofluorescence analysis, supporting a link to the CHS model. By controlling Tc17 generation, our data showed that NFATc1 acts a molecular switch in the development of Tc17 cells. Consequently, it controls the CD8+ T cell-mediated allergic skin inflammation. These findings will pave the way for future innovative therapeutic options for the treatment of allergic immune responses.



## Finite-time and fixed-time synchronization control of fuzzy Shunting Inhibitory Cellular Neural Network

Zhangir Nuriyev\*<sup>1</sup>

<sup>1</sup> Nazarbayev University

Email : zhangir.nuriyev@nu.edu.kz

### Abstract

Several problems based on finite-time synchronization are explored in this paper. The primary objective is to design the controllers for the various models based on Fuzzy Shunting Inhibitory Cellular Neural Networks (FSICNNs) and find out sufficient conditions for the systems' solutions to reach synchronization in finite time. In other words, we prove the existence of finite-time synchronization for three FSICNNs models that have not been studied before and suggest both the controllers and Lyapunov functions that would yield the feasible convergence time between solutions. Namely, we study models of delayed FSICNNs with and without inertial terms and FSICNNs with diffusion and without delays. Derived by means of the well-known in the scope of calculus maximum-value approach criteria, we give an upper bound to the time by which the synchronization phenomenon is guaranteed to occur in the three FSICNNs models. These results are supported by computer simulations showing the solutions' behavior for different initial conditions of FSICNNs.

# On the Hyers-Ulam stability and the Hyers-Ulam- Rassias stability of nonlinear iterative integro- differential equations with variable delays

Cemil Tunc\*<sup>1</sup> and Osman Tunc<sup>2</sup>

<sup>1</sup>Department of Mathematics Faculty of Sciences Van Yuzuncu Yil University 65080, Campus, Van, Turkiye

<sup>2</sup> Department of Computer Baskale Vocational School Van Yuzuncu Yil University 65080, Campus, Van, Turkiye

Email : cemtunc@yahoo.com

## Abstract

This work is interested in the existence and uniqueness of solutions, the Hyers-Ulam stability (HUS) and the Hyers-Ulam- Rassias stability (HURS) of an iterative integrofunctional differential equation (IIDDE) of first order with variable delays on a bounded interval, half infinite intervals and an infinite interval using the Banach fixed point theorem. The outcomes of this paper have scientific novelties and provide new contributions to the topic of the existence and uniqueness of solutions and the Ulam stabilities of IIDDEs.

**Keywords:** Uniqueness of solutions, Ulam stabilities, IIDDEs, fixed point theorem

## References

- [1] Egri, E., Ulam stabilities of a first order iterative functional-differential equation. *Fixed Point Theory* 12 (2011), no. 2, 321328.
- [2] Jung, S.-M., A fixed point approach to the stability of a Volterra integral equation. *Fixed Point Theory Appl.* 2007, Art. ID 57064, 9 pp.
- [3] Ulam, S.M., *Problems in modern mathematics*. Science Editions John Wiley & Sons, Inc., New York, 1964.
- [4] Graef, John R., Tun, C., engn, M., Tun, O., The stability of nonlinear delay integrodifferential equations in the sense of Hyers-Ulam. *Nonauton. Dyn. Syst.* 10 (2023), no. 1, (in press).

# New Results on Integro-Differential Equations with Variable Retardation

OSMAN TUNC<sup>\*1</sup> and CEMIL TUNC<sup>2</sup>

<sup>1</sup> Department of Computer Programing, Baskale Vocational School, Van Yuzuncu Yil University, Campus, 65080, Van, Turkey

<sup>2</sup> Department of Mathematics, Faculty of Sciences, Van Yuzuncu Yil University, 65080, Campus, Van Turkey

Email : osmantunc89@gmail.com

## Abstract

The paper is interested in a class of nonlinear time varying retarded integro- differential equations (RIDEs). By the functional Lyapunov-Krasovskii method, two new results having weaker conditions related to the uniformly stability (US), uniformly asymptotic stability (UAS), integrability, boundedness and boundedness at infinity of solutions of that RIDEs are given. We define a Lyapunov Krasovskii functional (LKF) for that system of RIDEs and perform the proof of the given results. Two examples are provided with their graphs to affirm the proposed theorems. The study of the results of this paper shows that the proposed theorems are not only applicable to the time varying linear RIDEs, but also applicable to the time varying nonlinear RIDEs.

## References

- [1] Pinelas, S., Tunc, O. Solution estimates and stability tests for nonlinear delay integro-differential equations. *Electron. J. Differential Equations* 2022, Paper No. 68, 12 pp.
- [2] Bohner, M., Tunc, O. Qualitative analysis of integro-differential equations with variable retardation. *Discrete Contin. Dyn. Syst. Ser. B* 27 (2022), no. 2, 639657.
- [3] Graef, J. R., Tunc, O. Asymptotic behavior of solutions of Volterra integro-differential equations with and without retardation. *J. Integral Equations Appl.* 33 (2021), no. 3, 289300.

# Investigation to Explore the Relationship Between Awareness and Behavior Change Towards NPI and Vaccine Usage in the Transmission of COVID-19 in Thailand

Alhassan Ibrahim<sup>1,2</sup>, Usa Wannasingha Humphries \*<sup>1</sup>, Parinya SA<sup>1</sup>, Isa Abdullahi Baba<sup>1,2</sup> and Rahat Zarin<sup>1</sup>

<sup>1</sup> Department of Mathematics, Faculty of Science, King Mongkuts University of Technology, Thonburi (KMUTT), 126 Pracha Uthit Road, Bang Mod, Thung Khru, Bangkok 10140, Thailand

<sup>2</sup> Department of Mathematical Sciences, Bayero University, Kano, Nigeria

Email : usa.wan@kmutt.ac.th

## Abstract

The novel coronavirus has caused a worldwide pandemic and emphasizes the need for mathematical modeling to understand the transmission dynamics of the virus and guide effective public health strategies. In this paper, we develop a compartmental model to study the transmission dynamics of COVID-19 in Thailand. Our model considers the vaccine population, divides the infected population into two groups symptomatic and asymptomatic, the impact of non-pharmaceutical interventions (NPIs), and awareness on reducing the infectiousness of the population. The model is fitted using real data from Thailand and we perform qualitative analysis to examine the stability of the equilibrium point and determine the conditions that lead to asymptomatic stability of the corresponding endemic equilibrium point. We also convert the model to the Atangana-Baleanu fractional order and carry out numerical simulations. We conduct sensitivity analysis on the models basic reproduction number to identify the parameters that have the greatest impact on reducing its value, including contact, recovery, contagiousness, and education outreach or campaign. Our analysis reveals that a combination of strategies, including increased education outreach and NPI usage, can effectively curb the spread of COVID-19 in Thailand. Our findings highlight the importance of adapting ongoing public health strategies to combat the spread of COVID-19. Our conclusion is that mathematical modeling can play a critical role in guiding effective public health interventions to mitigate the impact of pandemics such as COVID-19.

**Keywords:** COVID-19, Equilibrium points, Stability analysis, Atangana-Baleanu, Sensitivity analysis

# Modeling the effect of radio-chemotherapy on tumor cells interaction with optimal control and global sensitivity analysis

Arjun Kumar<sup>\*1</sup>, Balram Dubey<sup>1</sup> and Uma S Dubey<sup>2</sup>

<sup>1</sup> Department of Mathematics, BITS Pilani, Pilani Campus, Pilani, Rajasthan, 333031, India

<sup>2</sup>Department of Biological Sciences, BITS Pilani, Pilani Campus, Pilani, Rajasthan, 333031, India

Email : p20210039@pilani.bits-pilani.ac.in

## Abstract

Oncologists and applied mathematicians are interested in understanding the dynamics of cancer-immune interactions, mainly due to the unpredictable nature of tumour cell proliferation. In this regard, mathematical modelling offers a promising approach to comprehend this potentially harmful aspect of biology. This paper presents a novel dynamical model that incorporates the interactions between cancer cells, healthy tissue cells, and immune-stimulated cells in the environment of radio-chemotherapy treatment. We analyze the equilibria, identify conditions that lead to chaotic dynamics, and rigorously demonstrate the existence of chaos. Local stability analysis is performed, and we investigate transcritical, saddle-node, and Hopf bifurcations analytically and numerically. We derive the stability and direction conditions for periodic solutions. Furthermore, we examine an optimal control problem in the model that describes the dynamics of tumour-immune interactions, considering treatments such as radiotherapy and chemotherapy as control parameters. Our goal is to utilize optimal control theory to reduce the cost of radio chemotherapy, minimize the harmful effects of medications on the body, and mitigate the burden of cancer cells by maintaining a sufficient population of healthy cells. Cost-effectiveness analysis is employed to identify the most economical strategy for reducing the disease burden. Additionally, we conduct a Latin hypercube sampling-based uncertainty analysis to observe the impact of parameter uncertainties on tumour growth, followed by a sensitivity analysis. Through computer simulations, we demonstrate how the dynamical behaviour changes with variations in system parameters. The numerical results validate the analytical findings and illustrate that a multi-therapeutic treatment plan can effectively reduce tumour burden within a few days of drug administration

## Inverse Problems in Delay Differential Equation

Isyaku Idris\*<sup>1</sup>

<sup>1</sup> School of Mathematics and Statistics, Central South University, Changsha, China.

Email : isyakud@gmail.com

### Abstract

The aim of the research is to introduce concepts of direct problems, moment problems and inverse problems in delay differential equation in oscillating material i.e vibrating string and electric circuit by using finite difference scheme derived from Taylor series. We are to find the unknown properties of vibrating string which are usually given while solving direct problems.

**Keywords:** Inverse problems, direct problems, moment problems, delay differential equation, Oscillating system, electric circuit.

### References

- [1] N. RUDRAIAH, MATHEMATICAL MODELING FOR UNIFYING DIFFERENT BRANCHES OF SCIENCE, ENGINEERING AND TECHNOLOGY, National Research Institute for Applied Mathematics (NRIAM) No. 492/G, 7th Cross, 7th Block (West), Jayanagar Bangalore - 560 070.
- [2] B. JADAMBA, A. A. KHAN, M. SAMA, INVERSE PROBLEMS OF PARAMETER IDENTIFICATION IN PARTIAL DIFFERENTIAL EQUATIONS. Proceedings of the Satellite Conference of ICM 2010.
- [3] [3] Isyaku Idris, Eigenvalues Inverse Problems in Three Dimensional Rhotrices, American International Journal of Research in Formal, Applied & Natural Sciences 2018.
- [4] Abul Hasan Siddiqi, Mohamed Al-Lawati, Messaoud Boulbrachene, Modern Engineering Mathematics, by Taylor & Francis Group, LLC, 2018.
- [5] Dmitry Batenkov, Topics in Inverse Problems, Lecture notes. Tel Aviv University.

# Hybrid Crossover Dynamics of Immuno-Chemotherapy with Gene Therapy and Time-Delay Model: Numerical Treatments

S. M. AL-Mekhlafi<sup>1</sup>, M. M. Abou Hasan <sup>\*2</sup>, K. Udhayakumar<sup>3</sup> and F. A. Rihan<sup>4</sup>

<sup>1</sup>Sana'a University, Faculty of Education, Mathematics Department, Sana'a, Yemen

<sup>2</sup> Emirates aviation university

<sup>3</sup> Zayed University, UAE

<sup>4</sup>United Arab University, UAE

Email : eaus419@eau.ac.ae

## Abstract

Piecewise fractional differential equations (deterministic-stochastic differential equations or vice versa) have been introduced recently in literature. The purpose of this piecewise approach is to study effectively the model with real data. In this paper, a mathematical model of immuno-chemotherapy with gene therapy and the time delay is extended using the stochastic-deterministic piecewise hybrid fractional derivatives, where the hybrid fractional order and variable-order fractional operators are applied to extend the deterministic models. Two numerical methods are constructed to study the behavior of the proposed models. These methods are the nonstandard Caputo proportional constant finite difference method for solving the hybrid fractional deterministic models and the nonstandard Euler Maruyama technique to solve the stochastic model. Several numerical tests demonstrate the efficiency of the method and support the theoretical results.

# Investigation of The stability of The $F^*$ Iterative Algorithm on Strong Pseudocontractive Mappings And Its Applications

F. D. Ajibade<sup>\*1</sup>, O. O. Enoch<sup>1</sup> and T. P. Fajusigbe<sup>3</sup>

<sup>1</sup>Federal University Oye Ekiti, Department of Mathematics

2

Email : felix.ajibade@fuoye.edu.ng1

## Abstract

This paper is centered on conducting an inquiry into the stability of the  $F^*$  iterative algorithm to the fixed point of a strongly pseudo-contractive mapping in the framework of uniformly convex Banach spaces. To achieve the desired result, certain existing inequalities in convex Banach spaces were utilized as well as the stability criteria of Harder and Hicks. Other necessary conditions for the stability of the  $F^*$  algorithm on strong pseudo-contractive mapping were also obtained. Through a numerical approach, we prove that the  $F^*$  iterative algorithm is H-stable for strongly pseudo-contractive mapping. Finally, the solution of the mixed-type Volterra-Fredholm functional non linear integral equation is estimated using our results.

**Keywords and Phrases :** Stability,  $F^*$ -iterative algorithm, Pseudo-contractive mappings, nniformly convex Banach space, Mixed-type Volterra-Fredholm integral equation.



# Impulsive Delay Differential Equations with Delayed-Effectve Inputs

Jiaxu Li\*<sup>1</sup>

<sup>1</sup> University of Louisville

Email : [jiaxu.li@louisville.edu](mailto:jiaxu.li@louisville.edu)

## Abstract

It is well known that time delays exist in biological systems. Interventions are essential in many cases, such as drug administrations, fishery managements, vaccinations, etc. Current treatments for such interventions in mathematical modeling simulations and control theory are through impulse of perturbations of specific state variables and taking immediate effects, even though the effects of such interventions are delayed in most cases. To study the delayed effect of the biological interventions, we establish a novel concept for delay differential equation system with functional impulsive inputs that take effects in a future finite time interval. In this talk, we showcase this new modeling framework with the Hutchinson-Wright logistic population model, and the insulin administrations in artificial pancreas systems with a single pump (for insulin administration) or dual pumps (for both insulin and glucagon administrations).

# Bifurcation in a delayed predatorprey model with Holling type IV functional response incorporating hunting cooperation and fear effect

BENAMARA Ibtissam\*<sup>1</sup>

<sup>1</sup> Laboratory of Mathematics, Computer Science & Applications-Security of Information, Department of Mathematics, Faculty of Sciences, Mohammed V University in Rabat, street Ibn Battouta, 1014 Rabat, Morocco

Email : [ibtissam.benamara@um5.ac.ma](mailto:ibtissam.benamara@um5.ac.ma)

## Abstract

In this paper, we propose and analyze a delayed predatorprey model with Holling type IV functional response taking into account the cooperative hunting behavior in predators and the fear effect on prey due to predation risk. We first investigate the existence and stability of the model equilibria and show that the non-delayed model may lose its stability behavior via Hopf bifurcation by considering hunting cooperation as a bifurcation parameter. Next, we focus on the impact of fear on the dynamics of the delayed model and find that a Hopf bifurcation occurs for the delay as a bifurcation parameter. Furthermore, we numerically prove that the time-delay parameter may switch the stability behavior of the system. Before closing our study, numerical simulations are performed to illustrate our analytical findings.

## References

- [1] Alves MT, Hilker FM (2017) Hunting cooperation and Allee effects in predators. *J Theor Biol* 419:1322. <https://doi.org/10.1016/j.jtbi.2017.02.002>
- [2] Panday P, Samanta S, Pal N, Chattopadhyay J (2020) Delay induced multiple stability switch and chaos in a predatorprey model with fear effect. *Math Comput Simul* 172:134158
- [3] Hale JK, Lunel SMV (2013) *Introduction to functional differential equations*, vol 99. Springer

# Sensitivity Analysis and Optimal Control of Serial Killing

SHAMOONA JABEEN\*<sup>1</sup>

<sup>1</sup> Department of Mathematics, University of Science & Technology Bann-28100, Khyber Pakhtunkhwa, Pakistan

Email : shamoonajabeen@ustb.edu.pk

## Abstract

This work focuses on the elimination of Serial Killing by utilizing minimum resources. The initial crime transmission rate is examined using a mathematical model. We perform sensitivity analysis of the reproduction number  $\mathcal{R}$ , to find the most active parameters regarding the transmission of crime. These active parameters are intervened using suitable control variables. The objective function is designed to control the crime-addicted classes at a minimum cost. To minimize the objective functional Lagrangian and Hamiltonian are used. To check the validity of the results numerical simulations through MATLAB are performed.

## References

- [1] M. A. Mebratile, M. Y. Dawed, Mathematical Model Analysis of Crime Dynamics Incorporating Media Coverage and Police Force, *Journal of Mathematical and Computational Science*, (11):125-148, 2021.
- [2] K. S. Pritam, Sugandha, T. Mathur, S. Agarwal, Underlying dynamics of crime transmission with memory, *Chaos, Solitons, and Fractals*, (146):1-8, 2021.
- [3] P. Rakshit, S. Kumar, S. Noeiaghdam, et al., Modified SIR model for COVID-19 transmission dynamics: Simulation with a case study of UK, US and India, *Results in Physics*, 2022.
- [4] T.Padmavathi1, S. Senthamilselvi, S. S. Santra et al., Rotational Reaction over infected COVID-19 on human respiratory tract in the presence of sore effect with hall current. *The Bulletin of Irkutsk State University. Series Mathematics*, 40:1533, 2022.

## Analysis and stability of a social interaction model with age-structure and law enforcement

Syed Abbas<sup>\*1</sup>

<sup>1</sup> Indian Institute of Technology Mandi

Email : abbas@iitmandi.ac.in

### Abstract

In this work, we consider a social interaction model with age-structure and law enforcement. It is evident that age is an important factor in social dynamical models. We study stability of equilibrium solutions. This analysis is performed using basic reproduction number. Moreover, we study optimal control of considered problem. At the end, graphs are provided for illustration.

# Some Recent Studies Using Fractional Operators on the Incomplete Special Functions

NISHANT, SANJAY BHATTER\*<sup>1</sup>

<sup>1</sup> Department of Mathematics, Malviya National Institute of Technology Jaipur, India

Email : nishantjangra1996@gmail.com

## Abstract

Fractional calculus has numerous and varied applications in the domains of engineering and science, including optics, signals processing, viscoelasticity, fluid mechanics, electrochemistry, biological population models, and electromagnetics [2]. At this moment, the fractional calculus has opened its wings even larger to cover the dynamics of complex real world and new ideas are starting to be implemented and tested on real data. In order to better understand physical and engineering processes, fractional differential equations have been utilized to simulate them. In this study, the generalised fractional calculus operators will be examined and further formulate on the incomplete special functions [1]. The findings are presented in a concise manner that are helpful in creating certain lists of fractional calculus operators. A significant number of new and recognised outcomes including particularly the incomplete H-function and incomplete I-function follow as special instances of the primary discoveries.

## References

- [1] Jangid, K., Bhattar, S., Meena, S., Baleanu, D., Al Qurashi, M., & Purohit, S. D. (2020). Some fractional calculus findings associated with the incomplete I-functions, *Adv. Differ. Equ.*, 2020, 1-24.
- [2] Sun, H., Zhang, Y., Baleanu, D., Chen, W., & Chen, Y. (2018). A new collection of real world applications of fractional calculus in science and engineering. *Commun. Nonlinear Sci. Numer. Simul.*, 64, 213-231.

# A Seasonally Forced Eco-Epidemic Predator-Prey Model for the Impacts of Fear and Its Carry-Over Effect with Selective Predation

Sasanka Shekhar Maity <sup>\*1</sup>, Pankaj Kumar Tiwari<sup>2</sup> and Samares Pal<sup>1</sup>

<sup>1</sup> Department of Mathematics, University of Kalyani, Kalyani 741235, India

<sup>2</sup> Department of Basic Science and Humanities, Indian Institute of Information Technology, Bhagalpur 813210, India

Email : samaresp@yahoo.co.in

## Abstract

In many ecological situations, predation fear and its carry-over effect play crucial role in a predator-prey system. Predators have ability to distinguish prey that have been infected by some parasite, and have tendency to avoid the infected prey to reduce fitness cost. Here, we propose an eco-epidemic predator-prey model incorporating selective predation induced fear and its carryover effect. Also assume that the disease is of SI-type. In analytical part, we discuss basic properties like boundedness, feasible equilibria and its stability, and existence of Hopf bifurcation by taking carry-over effect as possible bifurcation parameter. Analytical findings are validated through numerical simulation. Our simulation results show that both the fear and its carry over effect destabilizes the system through limit cycle oscillations. Whereas, another fear factors: fear due to disease transmission and death of infected prey stabilizes the system by terminating the oscillations. To model the above situations more realistically, we consider the non-autonomous system by taking the parameters representing the cost of fear, carry-over effect, disease prevalence and selective preference of predators vary with time. For the seasonally forced system, conditions are obtained for which the system has at least one positive periodic solution. Our seasonally forced model showcases the appearance of a unique periodic solution, higher periodic solutions and complex bursting patterns

# New properties of the oscillatory behavior of neutral differential equations of higher order

Omar Bazighifan<sup>1</sup>

<sup>1</sup>Department of Mathematics, Faculty of Education, Seiyun University, Hadhramout, Yemen

Email : o.bazighifan@gmail.com

## Abstract

In this research, we applied three techniques: comparison technique, Riccati technique and integral averages technique to obtain several conditions and properties of the oscillatory behavior of neutral differential equations of higher-order. As proof of the effectiveness of new conditions and properties, we present some examples.

**Keywords:** Oscillation criteria, Higher order, Differential equations

# Optimal Control of an Influenza Transmission Model with Vaccination and Therapy

K. Arun Kumar<sup>\*1</sup>, A. Venkatesh<sup>1</sup>

<sup>1</sup> Department of Mathematics, AVVM Sri Pushpam College, (Affiliated to Bharathidasan University, Tiruchirapalli), Poondi, Thanjavur(Dt), Tamilnadu, India-613 503.

Email : arunkaruppusamyvinnai@gmail.com

## Abstract

The present work included the formulation and analysis of a non-linear epidemic mathematical model that incorporates the influence of vaccination and therapy in the context of Influenza transmission. This paper examines the stability of the disease-free equilibrium at both the local and global levels. Additionally, the model's fundamental reproduction number,  $\mathcal{R}_0$ , is calculated. The use of the normalised forward sensitivity index approach is being employed to ascertain the sensitivity analysis of the parameters in relation to the variable  $R_0$ . This study investigates the influence of vaccination and treatment rates on populations affected by infections. Two distinct types of controls are used to enhance the applicability of this model to an optimum control situation. The first control strategy aims to maintain the vaccination of persons who remain vulnerable, while the second control strategy seeks to enhance the treatment options available for affected individuals. The use of Pontryagin's Maximum Principle is employed for the purpose of analysing and numerically resolving the optimum control issue. Subsequently, the efficacy of the optimal control is evaluated by comparing the levels of the infected population with and without the implementation of controls. The integration of therapy and immunisation tactics has shown more favourable results, as it effectively reduces the prevalence of the illness within the designated period of control. **Keywords:** mathematical model, stability analysis, sensitive analysis, optimal control



# Multiple Periodic Solutions for Delay Differential Equations

FATIMA DIB <sup>\*1</sup> and NAIMA DAOUDI-MERZAGUI<sup>2</sup>

<sup>1</sup>Centre University of Maghnia, Algeria

<sup>2</sup> University of Tlemcen, Algeria

Email : fatimadib1967@yahoo.fr

## Abstract

In this work, we discuss the existence and the multiplicity of nonconstant subharmonic solutions for the following problem,

$$\begin{aligned}x''(t) + \lambda f(t, x(t), x(t - \tau)) &= 0, & t \in \mathbb{R}/\{t_j\}, \\-\Delta x'(t_j) &= I_j(x(t_j)), & j \in \mathbb{Z}, \\x(t) - x(t + 2r) &= x'(t) - x'(t + 2r) = 0,\end{aligned}\tag{5.1}$$

where  $r > 0$  and  $\lambda > 0$  are given constants,  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  is  $r$ -periodic in  $t$ .  $\Delta x'(t_j) = x'(t_j^+) - x'(t_j^-)$  with  $x'(t_j^\pm) = \lim_{t \rightarrow t_j^\pm} x'(t)$ ;  $t_j, j \in \mathbb{Z}$ , the instants where the impulses occur are such that  $0 = t_0 < t_1 < t_2 < \dots < t_p < t_{p+1} = 2r, t_{j+p+1} = t_j + 2r$ , for some fixed  $p \in \mathbb{N}$  and the function  $I_j : \mathbb{R} \rightarrow \mathbb{R}$  is continuous and satisfies  $I_{j+p+1} = I_j$ , for all  $j \in \mathbb{Z}$ . To study the solvability of (5.1), we use a variational approach based on an application of a variant of mountain-pass theorem.

## References

- [1] Bainov, D. and Simeonov, P., Systems with Impulse Effect, Ellis Horwood, Chichester, 1989.
- [2] Chen, H. and Li, J. Variational approach to impulsive differential equations with Dirichlet boundary conditions, Boundary Value Problems, Art. ID 324575, 16 pages, 2010.
- [3] Daoudi-Merzagui, N. and Dib, F. Positive periodic solutions to impulsive delay differential equations, Turkish Journal of Mathematics, Vol. 41, pp. 969-982, 2017.

## Optimal Control of a Periodically Switched Epidemic Model

SHRADDHA SALWAHAN\*<sup>1</sup>, SYED ABBAS<sup>1</sup> and ABDESSAMAD TRIDANE<sup>2</sup>

<sup>1</sup> School of Mathematical and Statistical Sciences,, Indian Institute of Technology Mandi,, Kamand (H.P.) - 175005, India

<sup>2</sup> Department of Mathematical Sciences, United Arab Emirates University,, Al Ain P.O. Box 15551, UAE

Email : d20075@students.iitmandi.ac.in

### Abstract

In this paper, we examine a modified Susceptible-Infected-Treatment-Recovered (SITR) model. The SITR model is assumed to be a periodically switched system. When  $\mathcal{R}_0$  is greater than 1, the control measures such as vaccination, isolation, and specific treatments are implemented. Given the limited availability of these resources, it is crucial to use them in the most efficient manner. The aim is to study the optimization of control strategies for infectious disease outbreaks in dynamic and resource-limited environments. In the first part of the article, we analyze the stability of the switched model and then formulate an optimal control problem of the periodically switched system. In the end, numerical simulations are performed to support the results discussed.

### References

- [1] Chowell G., & Brauer F., The basic reproduction number of infectious diseases: computation and estimation using compartmental epidemic models. *Mathematical and statistical estimation approaches in epidemiology*, 1-30, 2009.
- [2] Tridane A. Hajji M. A. & Mojica-Nava E., Optimal drug treatment in a simple pandemic switched system using polynomial approach. In *Mathematics Across Contemporary Sciences: AUS-ICMS, Sharjah, UAE, 227- 240*, 2017.
- [3] Liu X. & Stechlinski P., *Infectious disease modeling. A Hybrid System Approach*. Cham: Springer, 2017.

# Stability and Bifurcation Analysis of an SIR Epidemic Model

MEHMET GUMUS\*<sup>1</sup> and KEMAL TURK<sup>1</sup>

<sup>1</sup>Zonguldak Bulent Ecevit University, Faculty of Science, Department of Mathematics, 67100,  
Zonguldak, TURKEY

Email : m.gumus@beun.edu.tr

## Abstract

Throughout history, people have faced various epidemics, and the societal impacts caused by these epidemics have often been understood through mathematical modeling. Understanding and controlling the dynamics of disease spread represent fundamental challenges in the field of medical sciences. Mathematical epidemiology constitutes a significant component of research in this field. Especially in recent years, the outbreak of global pandemics like COVID-19 has further emphasized the importance of mathematical epidemiology. Research in this field plays a critical role in developing strategies to control and prevent epidemics. In this talk, we will consider a class of SIR epidemic model that divides the population into three classes, namely, susceptible, infected, and recovered individuals. First, we will demonstrate that the model is well-defined biologically. Thus, we will study dynamics analysis of the proposed model. We also anticipate that this model can be used to investigate the dynamics of various diseases. Finally, we will support our original and theoretical results with numerical examples.

## References

- [1] Brauer F, Castillo-Chavez C. *Mathematical models in population biology and epidemiology*. Vol. 2. New York: Springer; 2012.
- [2] Martcheva M. *An introduction to mathematical epidemiology*. Vol. 61. New York: Springer; 2015.
- [3] Kermack WO, McKendrick AG. A contribution to the mathematical theory of epidemics. *Proceedings of the Royal Society of London* 1927;A115(772):700-721.
- [4] Kaddar A. On the dynamics of a delayed SIR epidemic model with a modified saturated incidence rate. *Electronic Journal of Differential Equations* 2009;2009(133):1-7.

# On the Nonstandard Numerical Discretization of an SIR Epidemic Model

KEMAL TURK\*<sup>1</sup> and MEHMET GUMUS<sup>1</sup>

<sup>1</sup> Zonguldak Bulent Ecevit University, Faculty of Science, Department of Mathematics, 67100, Zonguldak, TURKEY

Email : kemalturk@beun.edu.tr

## Abstract

Epidemiological models are often formulated as systems of non-linear differential or difference equations. When a model represented by a nonlinear system of differential equations is given, this model can be discretized using certain methods. It is important to ensure that the discretized model retains as many dynamical properties of the continuous model as possible. For the purpose of discretization, Euler and Runge-Kutta methods, along with numerous other finitedifference techniques, are commonly used. In this talk, we will propose a class of discrete SIR model with vaccination. A nonstandard finite difference scheme (NSFD) will be developed. We will investigate the dynamical properties of discrete SIR epidemic model, including the nonnegativity and boundedness of solutions, the existence of equilibria, and the computation of the basic reproduction number. The local and global asymptotic stability of disease-free and endemic equilibria will be studied. Finally, some numerical simulations have been provided to illustrate our theoretical results.

## References

- [1] Allen LJS. Introduction to mathematical biology. New Jersey: Pearson/Prentice Hall; 2007.
- [2] Thieme HR. Mathematics in population biology. Vol. 1. New Jersey: Princeton University Press; 2018.
- [3] Mickens RE. Applications of nonstandard finite difference schemes. Singapore: World Scientific; 2000.
- [4] Kermack WO, McKendrick AG. A contribution to the mathematical theory of epidemics. Proceedings of the Royal Society of London 1927;A115(772):700-721.
- [5] Cui Q, Yang X, Zhang Q. An NSFD scheme for a class of SIR epidemic models with vaccination and treatment. Journal of Difference Equations and Applications 2014;20(3):416-422.
- [6] Suryanto A, Kusumawinahyu WM, Darti I, Yanti I. Dynamically consistent discrete epidemic model with modified saturated incidence rate. Computational and Applied Mathematics 2013;32:373-383.

## Numerical method for solving integro-differential model of biological species living together

Sondos M. Syam<sup>\*1</sup>, Z. Siri<sup>1</sup>, R. Md. Kasman<sup>1</sup>

<sup>1</sup>Institute of Mathematical Sciences, Universiti Malaya, 50603 Kuala Lumpur, Malaysia

Email: s2156213@siswa.um.edu.my

### Abstract

In this article, we investigate the solution of biological species living together model using the operational matrix method. This model consists of system of two fractional nonlinear integro-differential equations. We modify the operational matrix method to fit with this type of systems. We simplify the resulting algebraic system in practicable way. Instead of solving system of  $2m \times 2m$  equations, we simplify the problem to find the solution of  $2 \times 2$  of nonlinear equations. This for sure will reduce the computational cost when  $m$  is large. We proved the existence and uniqueness of the solution of biological species living together model. Also, we investigate two types of stability which are Ulam-Hyers stable and the generalized Ulam-Hyers stable. We implement the proposed method to solve three problems. We compare our results with the existence results in the literature when  $\alpha = 1$ . Also, we study the convergence of the solution when  $\alpha$  approaches to 1. The results show the efficiency of the proposed method.

# Problem with integral conditions for system of partial differential equations of third order

Grzegorz Kuduk<sup>1</sup>

<sup>1</sup>Faculty of Mathematics and Natural Sciences University of Rzeszow, Graduate of University, Rzeszow, Poland

E-mail: gkuduk@onet.eu

## Abstract

Let  $H(\mathbb{R}_+ \times \mathbb{R}^n)$  be a class of entire functions on  $\mathbb{R}$ ,  $K_L$  is a class of quasipolynomials of the form  $\varphi(x) = \sum_{i=1}^n Q_i(x) \exp[\alpha_i x]$ , where  $\alpha_i \in L \subseteq \mathbb{C}$ ,  $\alpha_k \neq \alpha_l$ , for  $k \neq l$ ,  $Q_i(x)$  are given polynomials.

Each quasipolynomial defines a differential operator  $\varphi\left(\frac{\partial}{\partial \lambda}\right)$  of finite order on the class of entire function, in the form

$$\sum_{i=1}^m Q_i\left(\frac{\partial}{\partial \lambda}\right) \exp\left[\alpha_i \frac{\partial}{\partial \lambda}\right] \Big|_{\lambda=0}.$$

In the strip  $\Omega = \{(t, x) \in \mathbb{R}^{n+1} : t \in ([T_1, T_2] \cup [T_3, T_4]), x \in \mathbb{R}^n\}$  we consider of the system of equations

$$\frac{\partial^3 U_i}{\partial t^3} + \sum_{j=1}^n \left\{ a_{ij} \left(\frac{\partial}{\partial x}\right) \frac{\partial^2}{\partial t^2} + b_{ij} \left(\frac{\partial}{\partial x}\right) \frac{\partial}{\partial t} + c_{ij} \left(\frac{\partial}{\partial x}\right) \right\} U_j(t, x) = 0, \quad (5.2)$$

$$\int_{T_1}^{T_2} e^{\alpha_k t} U_{ik}(t, x) dt + \int_{T_3}^{T_4} e^{\beta_k t} U_{ik}(t, x) dt = \varphi_{ik}(x), \quad k = 1, 2, 3 \quad (5.3)$$

where  $a_{ij}\left(\frac{\partial}{\partial x}\right)$ ,  $b_{ij}\left(\frac{\partial}{\partial x}\right)$ ,  $c_{ij}\left(\frac{\partial}{\partial x}\right)$ , for  $i = 1, \dots, n$ , are differential expression with entire symbols  $a_{ij}(\lambda) \neq 0$ ,  $b_{ij}(\lambda) \neq 0$ ,  $c_{ij}(\lambda) \neq 0$ ,  $\alpha, \beta \in \mathbb{R}$ . Let be  $\eta(\lambda) = \int_{T_1}^{T_2} W^{n-1}(t, \lambda) dt + \int_{T_3}^{T_4} W^{n-1}(t, \lambda) dt$  is a certain function  $W(t, \lambda)$  is a solution of equation  $\left(\frac{d^n}{dt^n} + \sum_{i=1}^n a_i(\lambda) \frac{d^{n-i}}{dt^{n-i}}\right) W(t, \lambda) = 0$ , satisfies conditions  $W^n(t, \lambda) \Big|_{t=0} = 1$ ,  $W^{n-1}(t, \lambda) \Big|_{t=0} = W(t, \lambda) \Big|_{t=0} = 0$ .

Denote be  $P = \left\{ \eta(\lambda) = 0, \lambda \in \mathbb{C} \right\}$  set zeros of function  $\eta(\lambda)$ .

**Theorem.** Let  $\varphi_{ki}(x) \in K_L$ ,  $i = 1, \dots, n$ ,  $k = 1, 2, 3$ ,  $j = 1, \dots, n$  then the class  $K_{L \setminus P}$  exist and unique solution of the problem (1) - (2). Solution of the problem (1) - (2) can be represented in the form

$$U_i(t, x) = \sum_{k=1}^3 \sum_{p=1}^n \varphi_{kp} \left(\frac{\partial}{\partial x}\right) \left\{ \frac{1}{\eta(\lambda)} T_{kjp}(t, \lambda) \exp[\lambda x] \right\} \Big|_{\lambda=0},$$

where  $T_{kjp}(t, \lambda) = l^T \left(\frac{d}{dt}, \lambda\right)$  is transpose of a matrix  $\left(\frac{d}{dt}, \lambda\right)$ .

Solution of the problem (1) - (2) according to the differential-symbol method [1] and [2] exists and unique in the class of quasi-polynomials.

## **References**

- [1] Kalenyuk P. I., Nytrebych Z. M. Generalized Scheme of Separation of Variables. Differential-Symbol Method. Publishing House of Lviv Polytechnic Natyonalny University, 2002. – 292 p. (in Ukrainian).
- [2] Kalenyuk P.I., Nytrebych Z.M., Kohut I.V., Kuduk G., Problem for nonhomogeneous evolution equation of second order with homogeneous integral conditions, *Math. Methods and Phys.- Mech. Polia.*, 58(2) (2015), 7–19.

## Problem with integral conditions for system of evolution equations of third order

Grzegorz Kuduk<sup>1</sup>

<sup>1</sup>Faculty of Mathematics and Natural Sciences University of Rzeszow, Graduate of University, Rzeszow, Poland

E-mail: gkuduk@onet.eu

### Abstract

Let  $B$  be a Banach space,  $A : B \rightarrow B$  linear operator, for this operator arbitrary powers  $A^n$ ,  $n = 2, 3, \dots$  be also defined in  $B$ , denote by  $x(\lambda)$  the eigenvector of the operator  $A$ , which corresponds to its eigenvalue  $\lambda \in \Lambda \subseteq \mathbb{C}$ .

We consider of the system of equations

$$\frac{d^3 U_i}{dt^3} + \sum_{j=1}^n \left\{ a_{j,i}(A) \frac{d^2}{dt^2} + b_{j,i}(A) \frac{d}{dt} + c_{j,i}(A) \right\} U_j(t) = 0, \quad (5.4)$$

$$\int_{T_1}^{T_2} t^k U_{i,k}(t) dt + \int_{T_3}^{T_4} t^k U_{i,k}(t) dt = \varphi_{i,k}, \quad i = 1, \dots, n, k = 1, 2, 3. \quad (5.5)$$

where  $a_{j,i}(A), b_{j,i}(A), c_{j,i}(A)$  are abstract operators, with entire symbols  $a_{j,i}(\lambda), b_{j,i}(\lambda), c_{j,i}(\lambda) \neq \text{const}$ , for  $\lambda \in \Lambda$ ,  $\varphi_{i,k} \in B$ ,  $T_1 > T_2 > T_3, T_4 >$ ,  $([T_1, T_2]) \cup ([T_3, T_4]) \subset \mathbb{R}$ .

Solution of the problem (1), (2), (3) according to the differential-symbol method [1] exists and unique in the class of quasi-polynomials.

Solution of the problem(1) -(2) according to the differential-symbol method [1] and [2] exists and unique in the class of quasi-polynomials.

## References

- [1] Kalenyuk P. I., Nytrebych Z. M. Generalized Scheme of Separation of Variables. Differential-Symbol Method. Publishing House of Lviv Polytechnic Natyonalny University, 2002. – 292 p. (in Ukrainian).
- [2] Kalenyuk P.I., Nytrebych Z.M., Kohut I.V., Kuduk G., Problem for nonhomogeneous evolution equation of second order with homogeneous integral conditions, Math. Methods and Phys.- Mech. Polia., 58(2) (2015), 7–19.



# Stability and Hopf bifurcation analysis in a discrete and distributed delayed neural network model with inertial coupling

BOOTAN RAHMAN\*<sup>1</sup>

<sup>1</sup> Mathematics Unit, School of Science and Engineering, University of Kurdistan Hewler (UKH), Erbil, Kurdistan Region, Iraq

Email : bootan.rahman@ukh.edu.krd

## Abstract

This talk presents the investigation of stability and Hopf bifurcation in a neural network model featuring discrete and distributed delays for a pair of neurons with inertial coupling. For a general distribution kernel, we identify a subset of amplitude death region. To delineate the complete region of amplitude death, we explore various delay distributions, including Dirac delta functions and gamma distributions. We establish stability criteria for the trivial steady state within parameter spaces encompassing neural interaction history and coupling strength. Our analysis reveals the occurrence of both Hopf and steadystate bifurcations as the steady state loses stability. Employing normal form theory and center manifold theory, we derive explicit formulas to examine the stability, direction, and other characteristics of periodic solutions emanating from the Hopf bifurcation points. To validate the theoretical findings, we provide a numerical illustration.

# Malaria dynamics with bimodality of incubation period in hosts in a seasonal environment

Attila Dénes\*<sup>1</sup> and Mahmoud A. Ibrahim<sup>2</sup> and Gergely Röst<sup>1</sup>

<sup>1</sup> National Laboratory of Health Security, Bolyai Institute, University of Szeged, Hungary

<sup>2</sup> University of Szeged, Hungary and Mansoura University, Egypt

Email : denesa@math.u-szeged.hu

## Abstract

To describe the bimodal distribution of the incubation time of *P. vivax* malaria in Korea corresponding to empirical observations, we present a periodic compartmental model of delay differential equations for malaria transmission dynamics with two distinct exposed classes in the human population and including time-dependent parameters for mosquito birth and death rates as well as biting rates. The short-term incubation period is modelled by exponential distribution, while the long-term incubation is assumed to be of fixed length. We identify the basic reproduction number as the spectral radius of a linear operator and show that it is a threshold parameter for the global dynamics of the model. We apply the model to data from South Korea. Joint work with Mahmoud A. Ibrahim and Gergely Rst.

# Explicit solutions to delay differential equations and first-passage times of delayed Wiener processes

MARIO LEFEBVRE <sup>\*1</sup> and IBRAHIM TOURE<sup>1</sup>

<sup>1</sup> Polytechnique Montreal, Montreal, Canada

Email : mlefevre@polymtl.ca

## Abstract

First, delay differential equations of the form

$$y'(t) = c_1 e^{c_2 t} y(c_0 t) + c_3 y(t)$$

and

$$y'(t) = c_0 + c_1 y^2(c_2 t),$$

where  $c_0, \dots, c_3$  are constants, are considered. Explicit and exact solutions to these equations are obtained. Then, the problem of computing the distribution of the first-passage time to a given constant for a Wiener diffusion process  $\{W(t), t \geq 0\}$  is treated in the case when there is a random delay  $T$  in the observation of  $W(t)$ . Explicit results are derived.

## References

- [1] Rangkuti, Y.M. and Noorani, M.S.M., The exact solution of delay differential equations using coupling variational iteration with Taylor series and small term. *Bulletin of Mathematics*, 4(1), 115, 2012.
- [2] Schwarz, W., On the convolution of inverse Gaussian and exponential random variables. *Communications in Statistics, Theory and Methods*, 31(12), 21132121, 2002.

# On delay differential equations: a robust fractional-order autoimmune disease application using wavelet frames

MUTAZ MOHAMMAD\*<sup>1</sup>

<sup>1</sup> Zayed University

Email : Mutaz.Mohammad@zu.ac.ae

## Abstract

In this presentation, we unveil an innovative mathematical model designed to enhance our understanding of tumor-immune interactions. Through the ingenious fusion of fractional order derivatives, time delays, and the enchanting world of wavelet frames, we delve deep into the intricate web of this complex biological system. The inclusion of a time delay ( $\tau > 0$ ) mirrors the real-world phenomenon, where the immune response to tumor growth unfolds gradually. Moreover, fractional derivatives introduce memory and hereditary effects, elevating the models predictive capabilities, especially within deterministic systems.

## References

- [1] Smith, J. (2018). Fractional calculus and the theory of distributions. *Anais da Academia Brasileira de Ciencias*, 90(2), 1843-1867.
- [2] Podlubny, I. (1999). *Fractional differential equations: An introduction to fractional derivatives, fractional differential equations, and methods of their solution and application* (Vol. 198). Academic press.
- [3] Mallat, S. (2008). *A wavelet tour of signal processing: The sparse way*. Academic press.
- [4] Dunn, G. P., Bruce, A. T., Ikeda, H., Old, L. J., & Schreiber, R. D. (2012). Cancer immunoediting: from immunosurveillance to tumor escape. *Nature immunology*, 3(11), 991-998.
- [5] Strang, G. (1996). The wavelet transform. *IEEE Transactions on Information theory*, 1(2), 255-261
- [6] Kilbas, A. A., Srivastava, H. M., & Trujillo, J. J. (2006). *Theory and Applications of Fractional Differential Equations*. Elsevier.
- [7] F.A. Rihan. *Delay Differential Equations and Applications to Biology*. Springer, 1st Ed., 2021, doi.org/10.1007/978-981-16-0626-7.

# Existence and exponential stability of a periodic solution of an infinite delay differential system with applications to Cohen-Grossberg neural network models

A. Elmwafy <sup>\*1</sup>, José J. Oliveira<sup>2</sup> and César M. Silva<sup>1</sup>

<sup>1</sup>University of Beira Interior, R. Marquês de vila e Bolama, Covilh, Portugal

<sup>2</sup>University of Minho, Campus de Gualtar, Portugal

Email : ahmed.elmwafy@ubi.pt

## Abstract

We investigate both the global exponential stability and the existence of a periodic solution of a general differential equation with unbounded distributed delays. The main stability criterion depends on the dominance of the non-delay terms over the delay terms. The criterion for the existence of a periodic solution is obtained with the application of the coincide degree theorem. We use the main results to get criteria for the existence and global exponential stability of periodic solutions of a generalized higher-order periodic Cohen-Grossberg neural network model with discrete-time varying delays and infinite distributed delays. Additionally, we provide a comparison with the results in the literature and a numerical simulation to illustrate the effectiveness of some of our results.

## References

- [1] Elmwafy A, Oliveira JJ, Silva CM. Generalized non-autonomous Cohen-Grossberg neural network model. arXiv preprint arXiv:2309.10824. 2023 Sep 4.

## A numerical solution of initial value problems by walsh polynomials approach

KROLY NAGY\*<sup>1</sup>

<sup>1</sup>Institute of Mathematics and Computer Sciences, Faculty of Informatics, Eszterhazy Krolly Cotholic University, Eger, Hungary

Email : nagy2.karoly@uni-eszterhazy.hu

### Abstract

In 1975 Chen and Hsiao gave the numerical solution of the initial value problems of systems of linear differential equations with constant coefficients by the Walsh polynomials approach [1]. In that period Chen and Hsiao wrote several papers in which they showed the applicability of their procedure in different fields of science [2, 3]. Chen and Hsiao did not deal with the analysis of the proposed numerical solution. These results were improved by Gt and Toledo for initial value problems of differential equations with variable coefficients on the interval  $[0, 1[$  and initial value  $\xi = 0$  [5, 4]. They proved the results by the tools of Walsh-Fourier analyses developed agily in the last twenty years [7]. It is easily seen that we can present a differential equation with variable coefficients where its solution is not determined in the point  $\xi = 0$ . In the present talk, we discuss the general case while  $\xi$  can take any arbitrary value in the interval  $[0, 1[$ . We show the existence and uniform convergence of the numerical solution, as well. Moreover, we proposed a faster multistep numerical method to obtain directly the values of the numerical solution without generating the Walsh functions [6].

## Quadratic stochastic processes associated with SIR models

TAIMUN QAISAR\*<sup>1</sup>

<sup>1</sup> UAE University, UAE

Email : 202090059@uaeu.ac.ae

### Abstract

Tracing back to Bernsteins work [1], quadratic stochastic operators continue to be a crucial tool the examination of dynamical properties and modeling across numerous fields. Among numerous others, the one we aim to discuss is its uses in network security. To curb the menace of network worms, several new systems were devised in the form of models [2]. The description of these models through quadratic stochastic operators creates new possibilities of investigating them. In-depth research is done on a number of evolution algebraic features related to network worm models like SIS-SIR [3] and SIS-SIR-PSIRD[4]. SIR model in discrete version is given by

$$\begin{aligned}x'_1 &= x_1^2 + (1 - \beta)x_1x_2 + x_1x_3 \\x'_2 &= (1 - \gamma)x_2^2 + (1 - \gamma - \beta)x_1x_2 + (1 - \gamma)x_2x_3 \\x'_3 &= \gamma x_2^2 + x_3^2 + \gamma x_1x_2 + x_1x_3 + (1 + \gamma)x_2x_3\end{aligned}$$

By means of above mentioned SIR model, we are going to construct quadratic stochastic processes and study their asymptotic behavior. Moreover associated differential equations are also derived

### References

- [1] Bernstein, S. (1942). Solution of a mathematical problem connected with the theory of heredity. *The Annals of Mathematical Statistics*, 13(1), 53-61.
- [2] Wang, Y., Wen, S., Xiang, Y., and Zhou, W. (2013). Modeling the propagation of worms in networks: A survey. *IEEE Communications Surveys and Tutorials*, 16(2), 942-960
- [3] Uygun Jamilov and Manuel Ladra (2022) Evolution algebras and dynamical systems of a worm propagation model, *Linear and Multilinear Algebra*, 70:19, 4097-4116, DOI: 10.1080/03081087.2020.1864264
- [4] Adilova, F., Jamilov, U., and Reinfelds, A. (2023). On discretetime models of network worm propagation generated by quadratic operators. *Mathematical Modelling and Analysis*, 28(2), 194217. <https://doi.org/10.3846/mma.2023.15999>

## Bifurcation analysis on the impact of novel TB vaccine

Dipo Aldila\*<sup>1</sup>

<sup>1</sup>Department of Mathematics, Universitas Indonesia, Depok 16424, Indonesia

Email : aldiladipo@sci.ui.ac.id

### Abstract

This study introduces a modeling framework that incorporates various elements in the complex transmission of TB, including the progression from initial infection (de novo) at varying speeds, relapse, reinfection at different rates, treatment failure, limitations in healthcare resources, and, notably, the introduction of the new M72/AS01E vaccine. The model is based on a system of ordinary differential equations that describe the dynamics of six different compartments within the human population over time. The model's basic reproduction number plays a pivotal role, serving as a measure of endemicity that guides the trajectories of the model. It can lead to either a forward or backward bifurcation depending on whether its value crosses the threshold of one. The specific conditions that determine which type of bifurcation occurs are found to be closely tied to the availability of treatment. In cases where the backward bifurcation occurs, the failure to meet the treatment threshold results in continued endemicity, even when stringent contact restrictions and effective vaccination efforts reduce the basic reproduction number below one. The study examines how the model responds to various individual parameters to determine effective disease control measures. The results show that, with the chosen parameter values, only forward bifurcations around critical points are observed, and these bifurcations are influenced by the gradient of the basic reproduction number in relation to the parameters. The findings highlight the importance of implementing stringent contact restrictions, reducing relapse rates through healthy lifestyle improvements that boost the immune system, increasing vaccination rates and their efficacy in containing the spread of the disease. Additionally, contact restrictions are identified as a significant factor that can lead to the creation of an endemic bubble between two adjacent Hopf points, resulting in a hysteresis loop. Within this bubble, there is a specific infection rate at which the trajectory's period significantly increases, leading to a slow movement around a saddle point before eventually transitioning to a one-time cycle and converging towards a stable state. Through a twoparameter continuation analysis involving contact restrictions and vaccination, the study identifies regimes where various modes of multistability occur, providing insights for the strategic allocation of resources and interventions at different levels of endemicity while avoiding counterproductive measures and entrapment in undesirable states.

**Keywords:** TB, M72/AS01E vaccine, reproduction number, bifurcation



# Existence and stability of solution for fractional wave equation of $\beta(x)$ -Laplace type with viscoelasticity

Mohamed Saadaoui\*<sup>1,2</sup>

<sup>1</sup> Faculty of Technology, Amaal Tleidji University Laboratoire de Mathématiques Pures et Appliquées (LMPA) P.O. Box 37, Laghouat 03000, Algeria

<sup>2</sup>Laboratoire d'Aide Multicritère à la Décision et Recherche Opérationnelle P.O. Box 32, EL ALIA, Bab Ezzouar, Alger, Algeria

Email : saadaoui16@gmail.com

## Abstract

In this work, we study the initial boundary value problem for a fractional viscoelastic equation of the  $\beta(x)$ -laplace type

$$u_{tt} + k(-\Delta)_p^{\beta(x)}u(x, t) + k_0(-\Delta)_p^{\alpha(x)}u_t - \int_0^t g(t-s)(-\Delta)_p^{\beta(x)}u(s)ds = k_1|u|^{q(x)-2}u,$$

First, we prove the existence of solution by using the Faedo-Galerkin approximations and under suitable conditions. Then, by applying an integral inequality due to Komornik, we obtain the stability result.

**Keywords:** Fractional wave equation,

$\beta(x)$ -Laplacian, the Gagliardo seminorm, the Galerkin method, Komornik integral inequ

# Higher order numerical scheme for singularly perturbed delay differential equations with small shift arising in bio science

AKHILA MARIYA REGAL<sup>1</sup> and S DINESH KUMAR\*<sup>1</sup>

<sup>1</sup> SCHOOL OF ADVANCED SCIENCES, DEPARTMENT OF MATHEMATICS, VELLORE INSTITUTE OF TECHNOLOGY, VELLORE, TAMIL NADU, INDIA

Email : mathdinesh005@gmail.com

## Abstract

In this paper, We propose a finite difference scheme to solve singularly perturbed delay differential equation of reaction-diffusion type with small shift arising in bio science. The scheme derived is almost second order accurate. To evaluate its effectiveness, several test examples with varying perturbation and delay parameter are considered. The proposed scheme is compared with other existing numerical schemes. Error estimate and the rate of convergence are shown in the table for proposed numerical scheme.

## References

- [1] Lange G Charles and Robert M. Miura., Singular perturbation analysis of boundary value problems for differential-difference equations, *SIAM Journal on Applied Mathematics*,42(3), (1982), 502531, 2000.
- [2] J. Mallet-Paret and R.D. Nussbaum, A differential-delay equations arising in optics and physiology, *SIAM J. Math. Anal.* 20, 249292, 1989.
- [3] Doolan, E.P., Miller, J.J. H., Schilders, W.H. A., *Uniform Numerical methods for problems with initial and boundary layers*, Boole Press, Dublin, 1980.
- [4] Chawla, M. M., A fourth-order tridiagonal finite difference method for general two-point boundary value problems with non-linear boundary conditions, *IMA Journal of Applied Mathematics (Institute of Mathematics and Its Applications)*, 22.1, 8989, 1978.

## Multivariate models for assessing the effects of nanoparticles on bacterial growth

Nhat-Thong Le<sup>1</sup>, Thi-Thu-Hoai Nguyen<sup>1</sup>, Ton-Duc Do<sup>2</sup> and Tri Thanh Pham<sup>\*3</sup>

<sup>1</sup> Research Center for Infectious Disease, International University - VNU-HCM, HCM city, Vietnam

<sup>2</sup>Department of Robotics and Mechatronics, School of Engineering and Digital Sciences (SEDS), Nazarbayev University, Republic of Kazakhstan

<sup>3</sup> Department of Biology, School of Sciences and Humanities, Nazarbayev University, Republic of Kazakhstan.

Email : tri.pham@nu.edu.kz

### Abstract

The emergence and rapid spread of multidrug-resistant (MDR) bacterial strains is a public health concern. This issue arises primarily from the misuse and abuse of antibiotics, fostering the development of antibiotic-resistant strains. Due to a growing number of nosocomial and community-acquired infections caused by MDR pathogens, nanoparticles (NPs) with enhanced antibacterial activity have recently been proposed as an alternative to antibiotics. However, the capacity to predict bacterial growth in response to NPs concentrations and time remains limited. In this study, we attempted to conduct a comparative analysis using six different mathematical models to quantify the effects of two different nanoparticles (NPs), namely protoporphyrin IX (PPIX) and sulfur and nitrogen co-doped carbon nanoparticles (SN-CNPs), on the growth of *Bacillus subtilis* (*B. subtilis*) and *Escherichia coli* (*E. coli*). To achieve this, we took measurements of bacterial growth for 24 hours at various NPs concentrations and assess how well the bacterial growth profiles fit the models. We observed that the Baranyi models provide the greatest fit for PPIX-treated experiments with 10 different concentrations, with the average  $R^2$  values of 0.925 and 0.986 for *B. subtilis* and *E. coli*, respectively. For SN-CNPs treatment, Logistic models showed the best match with a mean  $R^2$  of 0.898 for *B. subtilis*, while Gompertz models ( $R^2 = 0.928$ ) indicated the best fit for *E. coli*. Next, we constructed General Additive Models (GAM) under the assumption of a Gaussian distribution to evaluate the impacts of both NPs concentration and time on the growth. Our models indicated a strong correlation with  $R^2$  values of 0.917 and 0.991 for PPIX-treated *B. subtilis* and *E. coli*, respectively. As anticipated, bacteria treated with SN-CNPs also showed this remarkable correlation, with  $R^2$  values of 0.970 for *B. subtilis* and 0.971 for *E. coli*. In conclusion, this study provided mathematical models that accurately predict the effects of NPs quantity and time on bacterial growth rate.

**Keywords:** Bacterial growth, General Additive Models, mathematical models, multivariate models, protoporphyrin IX (PPIX), Sulfur and Nitrogen co-doped carbon nanoparticles (SN-CNPs),

# On the existence and stability of a viscoelastic Petrovsky equation with retard vary and a Logarithmic source term

Khedidja Abidi\*<sup>1</sup>

<sup>1</sup> Faculty of Science, Amaal Tleidji University Laboratoire de Mathematiques Pures et Appliquees (LMPA) P.O. Box 37, Laghouat 03000, Algeria .

Email : khadidjaabidi3@gmail.com

## Abstract

In this work, we consider a nonlinear wave equation of viscoelastic Dirichlet type under certain assumptions on the initial data

$$|u_t|^{s(x)}u_{tt}(x, t) - \Delta^2u(x, t) - \Delta u_{tt} - \sigma(t) \int_0^t g(t-s)\Delta^2u(s)ds + \mu_1h_1(u_t(x, t)) + \mu_2h_2(u_t(x, t - \tau(t))) + k_0|u_t|^{q(x)-1}; u_t = |u|^{p(x)-1} \ln |u|^k$$

First, we demonstrate the global existence of the weak solution of the considered problem using the Faedo-Galerkin method. Then, under certain preliminary results, we end this work by studying the asymptotic behavior of the solution by the energy method by constructing a suitable Lyapunov function.

**Keywords:** Asymptotic comportement, Faedo-Galerkin, Global existence, Viscoelastic.

## References

- [1] M. Daoulatli, I. Lasiecka, T. Doundykov; Uniform energy decay for a wave equation with partially supported nonlinear boundary dissipation without growth restrictions, *Discrete Contin. Dyn. Syst.*, 2 (2009), 67-95.
- [2] S. Kouemou-Patcheu, On a Global Solution and Asymptotic Behaviour for the Generalized Damped Extensible Beam Equation *J. Differential Equations*, 135 (1997), 299-314 .

# Hopf bifurcation in a predator-prey system with allee effect

BAYA LAADJAL\*<sup>1</sup>

<sup>1</sup> Laboratory of applied mathematics, University of Biskra, Algeria

Email : baya.laadjal@univ-biskra.dz

## Abstract

In this paper a predator-prey model with Allee effect associated with the prey species production and Evelev functional response is investigated for its dynamical behaviours, such as boundedness and local stability analysis. In the absence of time delay, the structure of equilibria and their local stability have been analyzed. In the presence of time delay, some sufficient conditions of the local stability of the positive equilibrium and the existence of Hopf bifurcation are discussed by analysing the corresponding characteristic equation, and the properties of Hopf bifurcation are given by using the normal form theory and centre manifold theorem. Some numerical simulations are performed to support our analytic results

## References

- [1] Biswas S., M. Saifuddin, S.K. Sasmal, S. Samanta, N. Pal, F. Ababneh, and J. Chattopadhyay, A delayed preypredator system with prey subject to the strong Allee effect and disease, *Nonlinear Dyn.* 84(3), 15691594, 2016.
- [2] Ma Z.P., H.F. Huo, and H. Xiang, Hopf bifurcation for a delayed predatorprey diffusion system with Dirichlet boundary condition, *Appl. Math. Comput.* 311, 118, 2017.

# Asymptotic Behavior of the Stokes Problem with Robin condition in a Domain with tiny Holes

Chafia Karek<sup>1</sup> and Amar Ould-Hammouda<sup>\*2</sup>

<sup>1</sup> Mathematics Department, Faculty of sciences, university 20th August 1955, Skikda, Algeria

<sup>2</sup>Laboratory of Physics Mathematics and Applications, ENS

Email : amar.ouldhammouda@g.ens-kouba.dz

## Abstract

We consider the Stokes problem in a domain  $\Omega_{\epsilon\sigma_1\sigma_2}$  of  $\mathbb{R}^N$ ,  $N \geq 3$   $\epsilon$ -periodically perforated by holes of size  $r(\epsilon)$ , with  $r(\epsilon)/\epsilon \rightarrow 0$  as  $\epsilon \rightarrow 0$  We are interested in the asymptotic behavior of the velocity and pressure of the fluid as  $\epsilon \rightarrow 0$  we use the periodic unfolding method introduced by Cioranescu, Damlaian and Griso [5] and [6] that allow us to consider a general geometric fram work and so to generalize the result of Cioranescu,D.,Donato, P.,and Horia Ene [4] and Anca Capatina, Horia Ene [3]

## References

- [1] A. Bensoussan, J.-L. Lions and G. Papanicolaou, *Asymptotic Analysis for Periodic Structures*, North Holland, Amsterdam, 1978.
- [2] Allaie, G. (1989) Homogenization of Navier-Stokes equation with a slip boundry condition. *Commun. Pure Appl. Math.* XLIV,605-642.
- [3] A.Capatina& H. Ene Homogenesation of the Stokes problem with a pure nonhomogeneous slip boundary contion by the periodic unfolding method. *Euro. Jnl of Mathematics* (2011), vol. 22,pp. 333-345
- [4] D.Cioranescu, P.Donato and H.Ene, Homogenisation of Stokes problem with non homogeneous slip boundary conditions; *Math.Meth. Appl. Sci.* 19(1996),857-881.
- [5] D. Cioranescu, A. Damlaian and G. Griso, Periodic unfolding and homogenization, *C. R. Acad. Sci. Paris, Ser. I*, 335 (2002), 99104.
- [6] D. Cioranescu, A. Damlaian and G. Griso, The periodic unfolding method in homogenization, to appear in *SIAM J. of Math. Anal.* (2008).

## Dynamics of hearing loss caused by noise hazard with time delay

KARMAND K. AHMAD<sup>\*1</sup>, GRACE O. AGABA<sup>2</sup> and BOOTAN RAHMAN<sup>3</sup>

<sup>1</sup> Department of Mathematics, Faculty of science, Soran University, Soran, Erbil, Kurdistan region, Iraq

<sup>2</sup> Department of Mathematics and Computer Science, Benue State University, Makurdi, Nigeria

<sup>3</sup> Department of Mathematics and Computer Science, Benue State University, Makurdi, Nigeria  
Email : karmand.ahmad@soran.edu.iq

### Abstract

Hearing loss, a growing public health concern primarily stemming from exposure to loud noise, is studied through a mathematical model employing delay differential equations. The model helps us grasp the dynamics of how noise impacts hearing and identify a critical noise level where the damage becomes unmanageable. Analytically, weve determined stability regions, demonstrating that the non-trivial steady state is stable below a critical time delay. Beyond this threshold, it becomes unstable and undergoes a Hopf bifurcation when the time delay matches the critical value. Our findings emphasize the need for comprehensive strategies to prevent and manage hearing loss, underscoring the significance of controlling noise pollution for an improved quality of life. Numerical simulations validate our analysis.

# Study the dynamic epidemiological model for dengue transmission with incubation period and saturated treatment function

PRASENJIT DAS\*<sup>1</sup>

<sup>1</sup>SRICHANDA MAHENDRANATH MEMORIAL INSTITUTION

Email : pdas1970@gmail.com

## **Abstract**

This article concentrates on the dynamical behaviour of Dengue infection model with multiple endemic equilibrium points consists of Susceptible, Infected and Recovered Class. Incubation period of the disease and saturation treatment function are incorporated with this model to study the effect of delay parameter on the transmission dynamics of the disease model along with the role of other parameters involved with this model particularly in connection with treatment functions are also investigated here to control the infection. Comprehensive numerical simulations are carried out to verify the result with biological interpretations.



# Reliability Modelling of Heterogeneous Data by Using Different Competing Weibull Mixture Models

Emad E. Elmahdy\*<sup>1</sup>

<sup>1</sup> Department of Mathematics, Science College, King Saud University, Riyadh 11451, P.O. 2455, Saudi Arabia

Email : Corresponding-author-eelmahdy@ksu.edu.sa

## Abstract

In this paper, the main focus of this study is to introduce different Weibull Mixture Distribution (WMD) models that are used for modelling life data and how to distinguish between them. A case study has been executed on a sample that includes both ordered exact times-to-failure and censoring (suspensions). The parameters estimates for WMD models are calculated by using the MLE method through the expectation-maximization (EM) algorithm. The goodness of fit tests are used for comparison of WMD models to determine the model of best fit to the data. From a mathematical statistics point of view, we found that modelling multimodal lifetime data under consideration with 3-fold WMD is the best choice.

**Keywords:** Life Data Analysis, Weibull Mixture Distribution (WMD), Competing Risk Weibull Mixture Distribution (CRWMD), Compound Competing Risk Weibull Mixture Distribution CCRWMD, Maximum Likelihood Estimation (MLE) Method, Expectation - Maximization (EM) Algorithm, Goodness of Fit (GOF) Tests, KolmogorovSmirnov (KS), The negative log-likelihood value  $-l(\theta)$ , The squared value for the correlation coefficient  $r^2$ .

## 6 List of Participants

Name	Email	Affiliation
Abbas, Syed	abbas@iitmandi.ac.in	Indian Institute of Technology Mandi ,India
Abidi, Khedidja	khadidjaabidi3@gmail.com	Amar Telidji University, Algeria
Ahmad, Karmand Khdr	karmand.ahmad@soran.edu.iq	Soran University, Iraq
Aldila, Dipo	aldiladipo@sci.ui.ac.id	<b>Universitas Indonesia, Indonesia</b>
Aldurayhim, Abdullah	Am.aldurayhim@psau.edu.sa	Prince Sattam bin Abdulaziz University, SA
Al-Salti, Nasser Said	alsalti@nu.edu.om	National University of Science and Technology, Oman
AlSariri, Tahani	alsariri@squ.edu.om	Sultan Qaboos University, Oman
Alshamsi, Eiman	eiman.alshamsi@uaeu.ac.ae	<b>UAE university, UAE</b>
Alsuraiheed, Turk	talsuraiheed@ksu.edu.sa	<b>King Saud University, SA</b>
AlKhabyah, Ali	Amalkhabyah@gmail.com	King Khalid University, SA
ALARYDAH, MOTASSEM	motassem.alarydah@ku.ac.ae	Khalifa University, UAE
Almaqrashi, Kifah	Kifahhsm@gmail.com	Sultan Qaboos University, Oman
Al-Musalhi, Fatma	fatma@squ.edu.om	Sultan Qaboos University, Oman
Almusawa, Musawa	malmusawi@jazanu.edu.sa	Jazan University, SA
Alsakaji, Hebatallah	heba.sakaji@uaeu.ac.ae	<b>UAE University, UAE</b>
Ahuod, Ahuod	oshokry@ub.edu.sa	University of Bisha, Saudi Arabia
Anwar, Mohamed Naim	mohamed.naem@puaa.edu.eg	Pharos University in Alexandria, Egypt
Asamoah, Joshua Kiddy	jkkasamoah@knust.edu.gh	University of Science and Technology Kumasi, Ghana
AZAMOV, ABDULLA	abdulla.azamov@gmail.com	Institute of Mathematics Uzbekistan, Uzbekistan
Baba, Isa Abdullahi	iababa.mth@buk.edu.ng	Bayero University Kano, Nigeria
Behl, Amanpreet	700039847@uaeu.ac.ae	UAE University, UAE
Bocharov, Gennady	bocharov@m.inm.ras.ru	Academy of Sciences, Moscow, Russia
Chinnathambi, Rajivganthi	rajivganthi.c@vit.ac.in	Vellore Institute of Technology, India
Das, Kalyan	daskalyan27@gmail.com	National Institute of Food Technology, India
DAS, PRASENJIT	pdas1970@gmail.com	MEMORIAL INSTITUTION, India
Denes,Attila	denesa@math.u-szeged.hu	University of Szeged, Hungary
El-Khatib, Youssef	youssef_elkhatib@uaeu.ac.ae	<b>UAE University, UAE</b>
Elmahdy, Emad	eelmahdy@ksu.edu.sa	<b>King Saud University, Saudi Arabia</b>
Elmojtaba, Ibrahim	elmojtaba@squ.edu.om	Sultan Qaboos University, Oman
Elmwafy, A	ahmed.elmwafy@ubi.pt	Universidade da Beira Interior, Portugal
FORREST, Omar	100059868@ku.ac.ae	KHALIFA UNIVERSITY, UAE
Gaipov, Abduzhappar	abduzhappar.gaipov@nu.edu.kz	Nazarbayev University, Kazkstan
Ganesan, Soundararajan	soundararajan.ganesan@nu.edu	<b>Nazarbayev University Kazakhstan, Kazakhstan</b>
Gm, Mehmet	m.gumus@beun.edu.tr	Zonguldak Blent Ecevit University, Turkey
GODHANIYA, MANOJKUMAR	mgodhaniya76@gmail.com	SAURASHTRA UNIVERSITY, India
Hashish, Adel	ahashish@uaeu.ac.ae	<b>UAEU, UAE</b>
Ivanov, Anatoli	aivanov@psu.edu	Pennsylvania State University, USA
Jamal, Sameerah	sameerah.jamal@wits.ac.za	University of the Witwatersrand, South Africa
Kandasamy, Udhayakumar	udhai512@gmail.com	Zayed University, UAE
Kashkynbayev, Ardak	ardak.kashkynbayev@nu.edu.kz	<b>Nazarbayev University, Kazkstan</b>
Kathiresan, S	kathiresan.sivakumar@nu.edu.kz	<b>Nazarbayev University, Kazkstan</b>
Khan, Gulfaraz	g_khan@uaeu.ac.ae	UAE University, UAE
Khan, Tahir	tahir_k@uaeu.ac.ae	<b>UAE University, UAE</b>
Kumar S, Dinesh	mathdinesh005@gmail.com	Velloree Institute of technology, Vellore
Laadjal, Baya	baya.laadjal@univ-biskra.dz	University of Biskra, Algeria

Name	Email	Affiliation
Li, Jiayu	jiayu.li@louisville.edu	University of Louisville, USA
Li, Qing	liqingliyang@126.com	shijiazhuang traditional chinese hospital, China
M, Manivel	manivelmani718@gmail.com	Bharathidasan University, India
MAJI, CHANDAN	chandanmaji.ju@gmail.com	Jagannath Kishore College, India
MATUSSE, AMERICO	ajmatusse@gmail.com	University of Pretoria South Africa
Meyerhans, Andreas	andreas.meyerhans@upf.edu	Universitat Pompeu Fabra, Spain
Merad, Ahcene	merad_ahcene@yahoo.fr	University of Oum El Bouaghi, Algeria
Mohamed, Mutaz	Mutaz.Mohammad@zu.ac.ae	Zayed University, UAE
Mondal, Priyanka	pm9732644@gmail.com	Vellore Institute of Technology, India
Muhammad, Khalid	k.muhammad@uaeu.ac.ae	<b>UAE University, UAE</b>
Mukherjee, Debasis	mukherjee1961@gmail.com	Thakurpukur Kolkata, India
Mumtaz, Kinza	19070011@lums.edu.pk	Lahore University of Man. Sci, Pakistan
N, Manoj	kingmanoj029@gmail.com	manonmaniam Sundaranar University, India
Nali, Ibrahim	ibrahim.nali@usmba.ac.ma	University of szeged, Hungary
NAGY, KROLY	nagy2.karoly@uni-eszterhazy.hu	Eszterhazy Karoly Catholic University, Hungary
Noor Aziz, Muhamad Hifzhudin	hifz_din@um.edu.my	<b>University of Malaya, Malaysia</b>
Nuriyev, Zhangir	zhangir.nuriyev@nu.edu.kz	<b>Nazarbayev University, Kazakhstan</b>
Otkel, Madina	madina.otkel@nu.edu.kz	<b>Nazarbayev University, Kazkstan</b>
OULD-HAMMOUDA, AMAR	amar.ouldhammouda@g.ens-kouba.dz	Ecole Normale Superieure Kouba, Algeria
Pal, Samares	samaresp@yahoo.co.in	Indian Institute of Information Technology, India
Pham, Tri	tri.pham@nu.edu.kz	<b>Nazarbayev University</b>
Pundeer, Naeem Ahmad	pundir.naeem@gmail.com	Jadavpur University, Kolkata, India
QAISAR, TAIMUN	202090059@uaeu.ac.ae	<b>UAE University, UAE</b>
Rababah, Abdallah	rababah@uaeu.ac.ae	UAE University, UAE
Radouane, Yafia	radouane.yafia@uit.ac.ma	Ibn Tofail University, Morocco
Rajan, Rakkiyappan	rakkigru@gmail.com	Bharathiar University, India
Raza, Ali	alleerazza786@gmail.com	University of Punjab, Lahore, Pakistan
Rihan, Fathalla	frihan@uaeu.ac.ae	<b>UAE University, UAE</b>
Saadaoui, Mohamed	saadaouik16@gmail.com	Laghouat university
Shaheen, Fouzia	201990173@uaeu.ac.ae	UAE university, UAE
Sheraz, Umar	21070003@lums.edu.pk	Lahore University of Man Sci, Pakistan
Shukur, Ali	shukur.math@gmail.com	University of Kufa, Iraq
Salwahan, Shraddha	ss14.salwahan@gmail.com	Indian Institute of Technology, India
Sarmad, Ghilmana	202190243@uaeu.ac.ae	UAE University, UAE
Singh, Varsha	vs4268736@gmail.com	University of Delhi, India
Sivakumar, kathiresan	kathiresan.sivakumar@nu.edu.kz	Nazarbayev University, Kazkstan
Shyamsunder	skumawatmath@gmail.com	Malviya National Institute of Technology, India
Sweilam, Nasser	nsweilam@sci.cu.edu.eg	Cairo University, Egypt
Syam, Sondos	s2156213@siswa.um.edu.my	<b>Universiti Malaya, Malaysia</b>
Tahir, Alhaji	altahir1ng@gmail.com	Modibbo Adama University Yola, NG
Teshome Tilahun, Getachew	gmgech@gmail.com	Haramaya, Ethibia
Tridane, Abdessamad	a-tridane@uaeu.ac.ae	UAE University
Turk, Kemal	kemalturk@beun.edu.tr	Zonguldak Blent Ecevit University, Turkey
Tursynkozha, Aisha	aisha.tursynkozha@nu.edu.kz	Nazarbayev University, Kazkstan
Volpert, Vitaly	volpert@math.univ-lyon1.fr	CNRS University Lyon, France